arXiv:1207.7203v1 [math.AP] 31 Jul 2012

EXTENSION PROBLEM AND FRACTIONAL OPERATORS: SEMIGROUPS AND WAVE EQUATIONS

JOSÉ E. GALÉ, PEDRO J. MIANA, AND PABLO RAÚL STINGA

Abstract. We extend results of Caffarelli–Silvestre and Stinga–Torrea regarding a characterization of fractional powers of differential operators via an extension problem. Our results apply to generators of integrated families of operators, in particular to infinitesimal generators of bounded C_0 semigroups and operators with purely imaginary symbol. We give integral representations to the extension problem in terms of solutions to the heat equation and the wave equation. On the way, a formula for fractional powers of generators of integrated semigroups is shown, that generalizes the well-known formula for C_0 semigroups. To get an approximation result for the solution to the extension problem, we give a theorem on the growth of perturbated tempered α-times integrated semigroups that could be of independent interest.

1. Introduction and main results

Motivated by the study of regularity properties of solutions to nonlinear equations involving the fractional Laplacian $(-\Delta)^\sigma$, $0<\sigma<1$, L. Caffarelli and L. Silvestre looked for solutions of the Bessel-type differential equation

$$\begin{cases} u_{yy}+\frac{1-2\sigma}{y}u_y=-\Delta u, & y>0,\\ u(0)=f, \end{cases} \tag{1.1}$$

where $f\in L^2(\mathbb{R}^n)$ and $u:[0,\infty)\to L^2(\mathbb{R}^n)$ is of class C^2. They interpreted the function u satisfying the equation above as the harmonic extension of f to a fractional dimension $2-2\sigma$. Then u is given by a Poisson type integral formula in terms of f. The remarkable property widely used in applications is that the nonlocal operator $(-\Delta)^\sigma$ acting on f in the domain of Δ is localized through u. More precisely, there exists a constant $c_\sigma<0$ such that

$$\lim_{y\to 0^+} y^{1-2\sigma}u_y(y)=c_\sigma(-\Delta)^\sigma f, \tag{1.2}$$

see [4]. For applications of the method see for instance [3] and [5].

Let us consider the extension problem (1.1) with Δ being replaced by a generic (closed) linear operator A. In [31], P. R. Stinga and J. L. Torrea gave a novel point of view to the extension problem by introducing the language of operator semigroup theory. This allowed them to get a general Poisson formula in terms of the semigroup e^{tA} and to get a Harnack inequality for the fractional harmonic oscillator. Moreover they analyzed the extension problem using Bessel functions. On the base of those tools they extended the Caffarelli–Silvestre theorem to a fairly general class of positive second order differential operators $D=-A$ on L^2-spaces in domains of $\mathbb{R}^n$. Such a class includes, for instance, the Laplacian in bounded domains and elliptic Schrödinger operators $D=-\operatorname{div}(A\nabla)+V$ with suitable potentials V. Subsequently, the approach to the extension problem given in [31] has been applied in [32] to get Harnack's inequalities for fractional powers of elliptic differential operators

---

Key words and phrases. Extension problem, fractional operator, Dirichlet-to-Neumann map, heat equation, wave equation, operator semigroup, integrated families.

2010 *Mathematics Subject Classification.* Primary: 35C15, 35K05, 35L05, 47D06, 47D09, 47D62. Secondary: 35R01, 35R03, 35J10, 35J70, 46J15, 46N20, 47A52, 26A33.

Research partially supported by Project MTM2010-16679, DGI-FEDER, of the MCYTS, Spain, and Project E-64, D.G. Aragón, Spain. The third author was partially supported by grant COLABORA 2010/01 from Planes Riojanos de I+D+I.

(both with discrete or continuous spectrum), and in [30] to obtain a boundary Harnack's inequality for the fractional Laplacian on the torus.

Nevertheless, the results of [4] and [31] do not seem to apply, totally or partially, to other important differential operators as, for example, those having purely imaginary symbol, in particular the free Schrödinger operator $A = i\Delta$, or those operators pointed out in Section 2 below. On the other hand, besides its theoretical significance (see [36] for instance), fractional powers of operators appear in many concrete settings. They occur, for instance, when considering fractional kinetics and anomalous transport [37], in fractional quantum mechanics [22] and fluid dynamics [5], and also in mathematical finance when modeling with Lévy processes [3]. Recently, the characterization (1.2) given in [4] has been used to show that the fractional Laplacian $(-\Delta)^\sigma$ coincides with a certain conformally covariant operator P_σ on the hiperbolic space $\mathbb{R}^n \times \mathbb{R}^+$ from which $\mathbb{R}^n$ is to be seen as its boundary; see [7].

Hence it sounds sensible to characterize fractional power operators $(-A)^\sigma$, in terms of solutions of local equations like in (1.2), for the widest possible class of operators A. In the present paper, we show that the semigroup structure revealed in [31], as underlying problem (1.1), can be subsumed in a more general framework which includes generators of α-times integrated semigroups and certain distribution semigroups. Working within that wider setting, we cover a wide range of important operators. Moreover, we are able to find a new expression for the solution to the extension problem that involves the solution to the associated wave equation.

To state the two main results we quickly introduce some notation and definitions. Let $\mathcal{B}(X)$ denote the Banach algebra of bounded operators on a Banach space $(X, \|\cdot\|)$. For $\alpha \geq 0$, let $(T_\alpha(t))_{t\geq 0}$ be a strongly continuous on $[0,\infty)$ family in $\mathcal{B}(X)$ such that $\sup_{t>0} t^{-\alpha}\|T_\alpha(t)\| < \infty$.

Suppose that there exists a (unique) closed and densely defined operator A on X, and such that $\lambda - A$ is invertible whenever $\Re\lambda > 0$, with resolvent function

$$(\lambda - A)^{-1} f = \lambda^\alpha \int_0^\infty e^{-\lambda t} T_\alpha(t) f\, dt, \quad \Re\lambda > 0,\ f \in X. \tag{1.3}$$

Then we say that $(T_\alpha(t))_{t\geq 0}$ is a *globally tempered* (or *tempered*, for short) *α-times integrated semigroup* in $\mathcal{B}(X)$, and that A is its *generator*. Analogously, if $(T_\alpha(t))_{t\geq 0}$ and A are as before but, instead of (1.3), they satisfy the relation

$$(\lambda^2 - A)^{-1} f = \lambda^{\alpha-1} \int_0^\infty e^{-\lambda t} T_\alpha(t) f\, dt, \quad \Re\lambda > 0,\ f \in X, \tag{1.4}$$

then we say that $(T_\alpha(t))_{t\geq 0}$ is a *tempered α-times integrated cosine family* in $\mathcal{B}(X)$ with *generator* A (see [12], for instance). Cosine families extend to $\mathbb{R}$ as even functions. Generators of integrated semigroups or integrated cosine families admit fractional powers $(-A)^\sigma$ in the Balakrishnan sense for every $0 < \sigma < 1$, see (4.5).

By W^α we denote the Weyl fractional derivative, see Section 3 for definitions. For real θ such that $0 \leq \theta \leq \pi/2$, set $S_\theta = \{z \in \mathbb{C} : |\arg z| < \theta\}$ if $\theta > 0$, and $S_0 = (0,\infty)$. Put $\mathbb{C}^+ := S_{\pi/2} = \{z \in \mathbb{C} : \Re z > 0\}$.

Theorem 1.1. *Fix $\alpha \geq 0$. Let A be the generator of a tempered α-times integrated semigroup $(T_\alpha(t))_{t\geq 0} \subseteq \mathcal{B}(X)$ and $0 < \sigma < 1$. Then a solution $u : S_{\pi/4} \to \mathcal{B}(X)$ to the (vector-valued) differential equation*

$$\begin{cases} u''(z) + \frac{1-2\sigma}{z} u'(z) = -Au(z), & z \in S_{\pi/4}; \\ \lim\limits_{z\to 0, z\in S_{\pi/4-\eta}} u(z) = f, & f \in X, 0 < \eta \leq \pi/4; \end{cases} \tag{1.5}$$

is given by

$$u(z) = \frac{z^{2\sigma}}{4^\sigma \Gamma(\sigma)} \int_0^\infty W^\alpha \left(\frac{e^{-z^2/(4t)}}{t^{1+\sigma}} \right) T_\alpha(t) f\, dt, \quad z \in S_{\pi/4}. \tag{1.6}$$

This solution is uniformly bounded on subsectors of $S_{\pi/4}$; that is, there exists $C > 0$ such that

$$\|u(z)\| < C\|f\|, \quad \text{for all } z \in S_\eta,$$

for every η such that $0 \leq \eta < \pi/4$.

Moreover, if $f \in \mathcal{D}(A)$, the domain of A, then

$$\lim_{z\to 0^+} \frac{u(z)-f}{z^{2\sigma}} = c_\sigma(-A)^\sigma f = \frac{1}{2\sigma}\lim_{z\to 0} z^{1-2\sigma}u'(z), \tag{1.7}$$

where $c_\sigma = 4^{-\sigma}\Gamma(-\sigma)\Gamma(\sigma)^{-1} < 0$ and both limits hold through proper subsectors of $S_{\pi/4}$.

Conversely, for $f \in \mathcal{D}(A)$ and $z \in S_{\pi/4}$, we have

$$u(z) = \lim_{\varepsilon\to 0^+} \frac{1}{\Gamma(\sigma)} \int_0^\infty W^\alpha\left(\frac{e^{-z^2/(4t)}e^{-\varepsilon t}}{t^{1-\sigma}}\right) T_\alpha(t)(-A)^\sigma f\,dt, \tag{1.8}$$

where the limit holds uniformly on compact subsets of $S_{\pi/4}$, and, for $f \in \mathcal{D}(A)$ and $z \in \overline{S}_{\pi/4}$,

$$u(z) = f + \frac{1}{\Gamma(\sigma)} \int_0^\infty W^\alpha\left(\frac{e^{-z^2/(4t)}-1}{t^{1-\sigma}}\right) T_\alpha(t)(-A)^\sigma f\,dt. \tag{1.9}$$

Remark 1.2. Formula (1.9) extends the solution u which is initially defined on the open sector $S_{\pi/4}$ to its closure, the closed sector $\overline{S}_{\pi/4}$.

Each of the equalities in (1.7) can be seen as a limit of localizations of the (in general) nonlocal operator $(-A)^\sigma$. The second one provides, in analogy to which is obtained in [4] and [31], the characterization of the fractional power $(-A)^\sigma$ as a correspondence from the Dirichlet boundary condition $f \in \mathcal{D}(A)$ to the Neumann-type boundary condition $\lim_{z\to 0^+} z^{1-2\sigma}u'(z)$.

As we mentioned before, we get a new formula for the solution u in terms of the solution to the wave equation. The following is the result we obtain if one replaces the integrated semigroup in Theorem 1.1 with a globally tempered integrated cosine family. This gives us more formulas to be added as solutions of the extension problem (1.1).

We will use the notation $\mathcal{C}_\alpha(t)$ replacing $T_\alpha(t)$ to refer to cosine families.

Theorem 1.3. *Let A be the generator of a tempered α-times integrated cosine function $(\mathcal{C}_\alpha(s))_{s\in\mathbb{R}} \subseteq \mathcal{B}(X)$. Then the extension problem (1.5) for A admits a solution $u(z)$ holomorphic in $z \in \mathbb{C}^+$ given by*

$$u(z) = d_\sigma \int_0^\infty W^\alpha\left(\frac{z^{2\sigma}}{(z^2+t^2)^{\sigma+1/2}}\right) \mathcal{C}_\alpha(t)f\,dt, \tag{1.10}$$

where $d_\sigma = \frac{2\Gamma(\sigma+1/2)}{\sqrt{\pi}\Gamma(\sigma)}$.

In addition, if $f \in \mathcal{D}(A)$ then the solution $u(z)$ can be alternatively written for $z \in \mathbb{C}^+$ as

$$u(z) = f + \kappa_\sigma \int_0^\infty W^\alpha\left((z^2+t^2)^{\sigma-1/2} - t^{2\sigma-1}\right) \mathcal{C}_\alpha(t)(-A)^\sigma f\,dt, \tag{1.11}$$

whenever $\sigma \neq 1/2$, for $\kappa_\sigma = \frac{2\Gamma(1/2-\sigma)}{4^\sigma\sqrt{\pi}\Gamma(\sigma)}$, or

$$u(z) = f + \frac{1}{\pi}\int_0^\infty W^\alpha\left(\operatorname{Log}\left(\frac{t^2}{z^2+t^2}\right)\right) \mathcal{C}_\alpha(t)(-A)^{1/2} f\,dt,$$

if $\sigma = 1/2$. Here Log *is the principal branch of the logarithm with argument in $[0, 2\pi)$, and the fractional powers appearing in the integrals are referred to this logarithm.*

In the classical case $A = \Delta$ on $\mathbb{R}^N$, the integrated cosine family in the statement above $v_\alpha(t) = \mathcal{C}_\alpha(t)f$ provides by derivation a mild solution to the wave equation with initial data f and null initial velocity

$$\begin{cases} v_\alpha''(t) - \Delta v_\alpha = 0, & t > 0, \\ v_\alpha(0) = f,\ v_\alpha'(0) = g. \end{cases}$$

The representation of the solution to the extension problem via the solution to the wave equation as in (1.10)-(1.11) is new. In the case of (1.1) it reads

$$u(x,y) = d_\sigma \int_0^\infty W^\alpha\left(\frac{y^{2\sigma}}{(y^2+t^2)^{\sigma+1/2}}\right) v_\alpha(x,t)\,dt,$$

for $\alpha > (N-1)|\frac{1}{2} - \frac{1}{p}|$, since this is the value of α for Δ to generate an integrated cosine family, see [12, Proposition 3.2].

Uniformly bounded C_0-semigroups are globally tempered with $\alpha = 0$. Then Theorem 1.1 extends in particular the results obtained in [4, Sections 2 and 3] and [31, Theorem 1.1] to all infinitesimal generators of bounded C_0-semigroups of Banach spaces. An important class of these semigroups are the so-called bounded analytic semigroups $(T(z))_{z \in S_\theta}$ on sectors of the complex plane. It is known that a closed operator A is the infinitesimal generator of such a semigroup $T(z)$ if and only if it is the generator of a tempered integrated cosine family. Hence, one can express the solution u of (1.1) for such an operator A by any of the formulas contained in Theorem 1.1 and Theorem 1.3.

As regards more differential operators other than those considered in [4] and [31], and to which Theorem 1.1 applies, one gets the differential operators A on $L^p(\mathbb{R}^n)$ whose symbols are of the form $iq(\xi)$, $\xi \in \mathbb{R}$, where q is a real elliptic polynomial or a C^∞ homogeneous function on $\mathbb{R}^n \setminus \{0\}$ such that $q(\xi) = 0$ implies $\xi = 0$. Among them, there are the operators $\partial^{2n-1}/\partial x^{2n-1}$, $n \in \mathbb{N}$, in particular the operator $\partial^3/\partial x^3$ associated with the Korteweg–de Vries equation [17], or operators of the form $A = iL$ for suitable Laplacians L on Riemannian manifolds or Lie groups [6, 8, 9].

For the convenience of prospective readers, we collect in Section 2 a more detailed list of the examples mentioned above, as well as some short proofs of corollaries deduced from Theorem 1.1. So we give there the proof of Theorem 1.3 as part of these consequences.

To prove Theorem 1.1 we use arguments which work in an abstract framework of Banach algebras and operator semigroup methods, so they avoid the dependency on the Fourier transform (and the Perron method) as in [4] or on L^2 spectral methods (and the use of Bessel functions) dealt with in [31]. First we check a variant of equation (1.5) on some specific scalar functions; see Lemma 3.1 (ii) and Lemma 3.4 (iii). One of these functions lies in a suitable Sobolev algebra $\mathcal{T}^{(\alpha)}(t^\alpha)$, which is a convolution Banach algebra defined by means of Weyl fractional derivatives W^α; see Section 3. Then, via bounded algebra homomorphisms with domain $\mathcal{T}^{(\alpha)}(t^\alpha)$ (which play the role of distribution semigroups), the properties of these functions are transferred to operator semigroups.

On the way a new formula for the Balakrishnan fractional power $(-A)^\sigma$ is given in Theorem 4.1 below in terms of the integrated semigroup $T_\alpha(t)$:

$$(-A)^\sigma f = \frac{\Gamma(\sigma+\alpha+1)}{\Gamma(-\sigma)\Gamma(1+\sigma)} \int_0^\infty \left(T_\alpha(t) f - \frac{t^\alpha}{\Gamma(\alpha+1)} f \right) \frac{dt}{t^{\sigma+\alpha+1}}.$$

This formula is to be used in the proof of Theorem 1.1. We also establish in Theorem 4.2 a result on additive perturbations with the identity $A - \varepsilon$, $\varepsilon > 0$, of generators A, which might be of independent interest. Here, it is used to give a corollary about convergence of solutions u_ε of equation (1.5) for perturbations $A - \varepsilon$. This corollary is proved in Section 2. Other perturbation properties of generators of integrated semigroups can be found in [35].

Section 3 collects some preliminary results about functions in convolution Sobolev algebras. Section 4 contains, besides the proof of Theorem 4.1, the perturbation theory for globally tempered integrated semigroups. Finally, in Section 5 we prove Theorem 1.1.

Throughout the paper we assume the constant convention; that is, the same letter, say C, usually taken through the text as playing the role of a constant, may well have different values from a line to another.

2. Applications and related results

We start with a direct consequence of Theorem 1.1 and the results on perturbations of Section 4.

2.1. Approximation by perturbations.

Let A be the generator of a globally tempered α-times integrated semigroup $T_\alpha(t)$ and put $A_\varepsilon := A - \varepsilon$ for $\varepsilon > 0$. By Theorem 4.2 below, A_ε is also the generator of a globally tempered α-times integrated semigroup $T_{\alpha,\varepsilon}(t)$. We have the following result about approximation of solutions of equation (1.5).

Corollary 2.1. *Let u and u_ε be the solutions of problem (1.5) which correspond to the semigroups $T_\alpha(t)$ and $T_{\alpha,\varepsilon}(t)$ respectively. Then*

$$\lim_{\varepsilon\to 0^+} u_\varepsilon(z) = u(z)$$

in the norm of X and uniformly in z running over compact subsets of the sector $S_{\pi/4}$.

Proof. From the formula for $T_{\alpha,\varepsilon}(t)$ established in Theorem 4.2 one has that $\lim_{\varepsilon\to 0} T_{\alpha,\varepsilon}(t)f = T_\alpha(t)f$ for every $t > 0$, $f \in X$. On the other hand, if K is a compact subset of $S_{\pi/4}$ then one has that the function Φ_K defined by $\Phi_K(t) := \sup_{z\in K} |W^\alpha(e^{-z^2/4t}/t^{1+\sigma})|$ is integrable on $(0,\infty)$. Then it is enough to apply dominated convergence in formula (1.6) for $T_{\alpha,\varepsilon}(t)$ to get the result. □

2.2. **Bounded C_0-semigroups.** As pointed out before, Theorem 1.1 is also valid for infinitesimal generators A of uniformly bounded C_0-semigroups. In this case such semigroups can be seen as the solution to the heat equation for A,

$$\left\{\begin{array}{ll} w'(s) = Aw(s), & s>0; \\ w(0) = f, & f \in X. \end{array}\right. \tag{2.1}$$

The result reads as follows.

Theorem 2.2. *Let A be the generator of a bounded C_0-semigroup $(T(t))_{t\geq 0} \subseteq \mathcal{B}(X)$. Fix $0 < \sigma < 1$. Then a solution $u : S_{\pi/4} \to \mathcal{B}(X)$ to the differential equation*

$$\left\{\begin{array}{ll} u''(z) + \frac{1-2\sigma}{z} u'(z) = -Au(z), & z \in S_{\pi/4}; \\ \lim\limits_{z\to 0, z\in S_{\pi/4-\eta}} u(z) = f, & f \in X, 0 < \eta \leq \pi/4; \end{array}\right.$$

is given by

$$u(z) = \frac{z^{2\sigma}}{4^\sigma \Gamma(\sigma)} \int_0^\infty \frac{e^{-z^2/(4t)}}{t^{1+\sigma}} T(t)f\, dt, \quad z \in S_{\pi/4}. \tag{2.2}$$

This solution is uniformly bounded on subsectors of $S_{\pi/4}$.

Moreover, if $f \in \mathcal{D}(A)$ then one can express the fractional operator $(-A)^\sigma$ by

$$\lim_{z\to 0} \frac{u(z) - f}{z^{2\sigma}} = c_\sigma (-A)^\sigma f = \frac{1}{2\sigma} \lim_{z\to 0} z^{1-2\sigma} u'(z), \tag{2.3}$$

where $c_\sigma = 4^{-\sigma}\Gamma(-\sigma)\Gamma(\sigma)^{-1} < 0$ and both limits hold through proper subsectors of $S_{\pi/4}$.

Conversely, for $z \in S_{\pi/4}$ and $f \in \mathcal{D}(A)$, we have

$$u(z) = \lim_{\varepsilon\to 0^+} \int_0^\infty \frac{e^{-z^2/(4t)} e^{-\varepsilon t}}{t^{1-\sigma}} T(t)(-A)^\sigma f \frac{dt}{\Gamma(\sigma)}, \tag{2.4}$$

where the limit holds uniformly on compact subsets of $S_{\pi/4}$, and, for $f \in \mathcal{D}(A)$ and $z \in \overline{S}_{\pi/4}$,

$$u(z) = f + \frac{1}{\Gamma(\sigma)} \int_0^\infty \frac{e^{-z^2/(4t)} - 1}{t^{1-\sigma}} T(t)(-A)^\sigma f\, dt. \tag{2.5}$$

Formulas (2.2), (2.3) appear in [31, Theorem 1.1], where they are obtained for generators of semigroups arising as differential operators acting on $L^2(\Omega)$, for Ω open subset of $\mathbb{R}^N$. Theorem 2.2 above is valid for *all* infinitesimal generators of bounded C_0-semigroups on Banach spaces, so it supplies the widest possible extension of [31, Theorem 1.1] in the setting of operator semigroups; in particular it applies to (the suitable ones) differential operators on $L^p(\Omega)$, $1 \leq p \leq \infty$. The proofs of (2.2) and (2.3) given here, in Section 5 below, are different from those given in [31, Theorem 1.1] and do not rely on the spectral theorem or the Fourier transform.

Formulas (2.4) and (2.5) are new. The weight $e^{-\varepsilon t}$ and corresponding limit in (2.4) must be included because the vector-valued mapping $t \mapsto e^{-z^2/4t} t^{\sigma-1} T(t)(-A)^\sigma f$ is not necessarily Bochner-integrable on $(0,\infty)$ in general. When the space X is assumed to be reflexive and A is a (generally unbounded) spectral operator of scalar type (see [10] for the basic theory and L^∞-functional calculus of these operators), the weight and limit can be removed from (2.4). In fact, we have the following

Proposition 2.3. *Let X be a reflexive Banach space. Suppose that A is a spectral operator of scalar type with spectrum in $[0,\infty)$ and spectral measure $E(\lambda)$. Let $(T(z))_{\Re z>0}$ denote the holomorphic semigroup in $\mathcal{B}(X)$ generated by A. Then there exists a solution of the boundary value problem*

$$\begin{cases} u''(z)+\frac{1-2\sigma}{z}\,u'(z)=-Au(z), & \text{for } z\in S_{\pi/4};\\ \lim\limits_{z\to 0,z\in S_{\pi/4-\eta}} u(z)=f, & \text{for } f\in X,\ 0<\eta\leq \frac{\pi}{4}; \end{cases}$$

which satisfies (2.2),(2.3),(2.5) of Theorem 2.2 and, furthermore,

$$u(z)=\frac{1}{\Gamma(\sigma)}\int_0^\infty \frac{e^{-z^2/(4t)}}{t^{1-\sigma}}T(t)(-A)^\sigma f\,dt,\quad z\in S_{\pi/4},\ f\in\mathcal{D}(A).$$

Proof. By the L^∞-functional calculus for A (see [10, Theorem 11 c)] one can write

$$T(t)(-A)^\sigma f=\int_0^\infty e^{-t\lambda}\lambda^\sigma dE(\lambda)f,\quad f\in X,$$

so that for every ϕ in the dual space X' we have

$$\phi(T(t)(-A)^\sigma f)=\int_0^\infty e^{-t\lambda}\lambda^\sigma dE_{f,\phi}(\lambda),\quad f\in X.$$

Here $E_{f,\phi}(\lambda)$ is the complex measure defined by $E_{f,\phi}(\lambda):=\phi(E(\lambda)f)$. Then, for $a=\Re(z^2)$,

$$\begin{aligned}\int_0^\infty \frac{e^{-a/4t}}{t^{1-\sigma}}\left|\phi\left(T(t)(-A)^\sigma f\right)\right|dt &\leq \int_0^\infty\int_0^\infty \frac{e^{-a/4t}}{t^{1-\sigma}}\lambda^\sigma e^{-t\lambda}dt\,d|E_{f,\phi}(\lambda)|\\ &\leq \int_0^\infty\int_0^\infty r^{\sigma-1}e^{-r}dr\,d|E_{f,\phi}(\lambda)|\\ &=\Gamma(\sigma)\int_0^\infty d|E_{f,\phi}(\lambda)|\leq C\|f\|\|\phi\|.\end{aligned}$$

It follows that the integral of the statement is absolutely convergent and therefore we can apply the dominated convergence theorem in (2.4) to obtain the result. □

The result above applies to every self-adjoint closed operator on a Hilbert space, so in particular to non-negative self-adjoint differential operators on L^2 spaces. Hence Proposition 2.3 extends formula (1.7) of [31, Theorem 1.1].

2.2.1. *The case of analytic semigroups.* Many interesting examples of C_0-semigroups turn out to be analytic and bounded in sectors.

Definition 2.4. Let $\theta\in(0,\pi/2)$. An analytic semigroup $(T(z))_{z\in S_\theta}\subseteq\mathcal{B}(X)$ on S_θ is said to be bounded if $\lim_{z\to 0}T(z)f=f$ and $\|T(z)\|$ is bounded on every proper subsector of S_θ.

Remark 2.5. As a matter of fact, one has that if A generates a uniformly bounded C_0-semigroup then the fractional power operator $-(-A)^\rho$ is the infinitesimal generator of a bounded analytic semigroup for $0<\rho<1$; see [1, p. 238] or [36, p. 263].

Next, a collection of examples of bounded analytic semigroups is listed for further reference. All these examples arise in a common setting. We assume that (M,d,μ) is a metric measure space where M has regular volume growth (that is, M satisfies the double volume property), see [6], or, more generally, the volume of balls in M is of polynomial growth in radii at 0 and at infinity, see [8, 9]. Additionally, it is usually assumed that, for some open subset Ω of M, L is a non-negative self-adjoint operator on $L^2(\Omega,\mu)$ such that the semigroup e^{-tL} has an integral kernel p_t which satisfies a pointwise *Gaussian upper estimate* of order 2 [8] or, more generally, of order $m\geq 2$ [9, 6].

Under the type of conditions above there exists a real number $\tau\geq 0$ such that the semigroup $e^{-t(\tau+L)}$ admits an extension as a bounded analytic semigroup in $\mathbb{C}^+$ from $L^p(\Omega)$ into itself. Note that this means that for some $\nu\geq 0$ and a constant $C_{\nu,\tau}$ the property

$$\|e^{-zL}\|\leq C_{\nu,\tau}e^{\tau\Re z}\left(\frac{|z|}{\Re z}\right)^\nu,\quad z\in\mathbb{C}^+, \tag{2.6}$$

is fulfilled for the L^p norm, in this case. See [8, Lemma 2], [9, Theorem 20 and 25], [6, Theorem 4.3].

Example 2.6. Among the infinitesimal generators of semigroups which satisfy property (2.6) one has the following. Note that according to Remark 2.5 every fractional power, in particular the square roots, of any of such examples also satisfies (2.6), though it may well be with different exponents ν, τ.

(i) The Laplacian $-\Delta$ or, more generally, Schrödinger operators $L := -\Delta + V$ with V in the Kato class, and magnetic Schrödinger operators acting initially on $L^2(\mathbb{R}^N)$ [8, p. 178], [6, p. 303].
(ii) Laplace-Beltrami operators acting on complete Riemannian manifolds of bounded geometry or with non-negative Ricci curvature [8, p. 178], [6, pp. 299, 302].
(iii) Uniformly elliptic differential operators, of second or higher order, with measurable coefficients acting initially on $L^2(\Omega)$ for $\Omega = \mathbb{R}^N$ or a compact Riemannian manifold without boundary [8, p. 178], [9, pp. 159, 162, 165].
(iv) Bochner Laplacians acting on sections of vector bundles [8, p. 178].
(v) Sub-Laplacians acting on Lie groups of polynomial growth and stratified nilpotent Lie groups; see [6, p. 299] and references therein.
(vi) Distinguished sub-Laplacians acting on the Heisenberg group [27], [28], and generalized Heisenberg groups [29, Proposition 2.4.1].

Remark 2.7. As said before, in the examples above the semigroups are given by integral operators with corresponding integral heat kernels p_t. By writing down these kernels into formula (1.6) and applying Fubini's theorem we obtain Poisson formulas for the solution of the extension problem related to the fractional powers of the generators of these semigroups. In particular, if we do so with the Laplacian in $\mathbb{R}^n$ then we recover the Poisson formula given in [4]; see also [31].

Bounded analytic semigroups not only supply examples to which Theorem 1.1 applies. They also yield interesting integrated families of operators. We will deal with these cases later on.

2.2.2. *The extension problem and the wave equation.* Let A be a closed and densely defined operator on a Banach space X. The wave equation for A with null initial velocity is

$$\begin{cases} w''(s) = Aw(s), & s>0; \\ w(0) = f, & f \in \mathcal{D}(A); \\ w'(0) = 0; \end{cases} \tag{2.7}$$

It is known that equation (2.7) admits a unique classical solution if and only if A is the generator of a cosine function $\mathcal{C}_0(s)$, with $w(s) = \mathcal{C}_0(s)f$ [16, Theorem 8.2]. When A generates a ν-times integrated cosine function $\mathcal{C}_\nu(s)$ with $\nu > 0$ then it is still possible to find mild solutions to problem (2.7); see [2].

The following result is implicitly contained in [12] and [23] altogether. For the sake of completeness we outline a proof of it.

Lemma 2.8. *The operator A generates a globally temperated integrated cosine function if and only if A is the infinitesimal generator of a bounded analytic semigroup on $\mathbb{C}^+$.*

Proof. Suppose that A is the generator of an α-times integrated cosine function $\mathcal{C}_\alpha$ such that $\|\mathcal{C}_\alpha(s)\| \le C|s|^\alpha$, $s \in \mathbb{R}$. Then the operator $T(z)$ in $\mathcal{B}(X)$ defined by

$$T(z)f := \int_0^\infty W^\alpha \left(\frac{e^{-s^2/4z}}{\sqrt{\pi z}} \right) \mathcal{C}_\alpha(s)f\, ds, \quad f \in X, z \in \mathbb{C}^+, \tag{2.8}$$

is a holomorphic semigroup generated by A and such that $\|T(z)\| \le C(|z|/\Re z)^{\alpha+(1/2)}$, for all $z \in \mathbb{C}^+$. (This fact is shown in [23, p. 142] for positive integer α; for any fractional α the proof is similar).

Conversely, suppose now that A is the infinitesimal generator of an analytic semigroup $(e^{zA})_{\Re z>0}$ satisfying $\|e^{zA}\| \le C(|z|/\Re z)^\alpha$, for all $z \in \mathbb{C}^+$ and some $\alpha \ge 0$. Then $-\sqrt{-A}$ generates a holomorphic semigroup such that $\|e^{-z\sqrt{-A}}\| \le C(|z|/\Re z)^\beta$, for all $z \in \mathbb{C}^+$ and $\beta = \alpha+(1/2)$; see [12, Theorem 4.1], [15, pp. 342, 343]. Hence $i\sqrt{-A}$ generates a β-times integrated group $T_\beta(it)$ with $\|T_\beta(it)\| \le C|t|^\beta$,

$t \in \mathbb{R}$ (note Theorem 2.9 below). Finally, it follows from [12, Lemma 3.1] that $A = (i\sqrt{-A})^2$ generates a β-times integrated cosine function $\mathcal{C}_\beta := (T_\beta(it) + T_\beta(-it))/2$, which obviously satisfies $\|\mathcal{C}_\beta(t)\| \leq C|t|^\beta$, for all $t \in \mathbb{R}$. □

We next proceed to give the proof of Theorem 1.3. So we assume that A is the generator of an integrated cosine function $\mathcal{C}_\alpha(s)$ such that $\|\mathcal{C}_\alpha(s)\| \leq C|s|^\alpha$, $s \in \mathbb{R}$. Notice that, according to the argument employed in the preceding lemma, the exponent above would be $\alpha + 1/2$ if we had started assuming that A is the generator of a bounded analytic semigroup as the one in (2.6) with $\nu = \alpha$.

Proof of Theorem 1.3. Substituting the expression of $T(t)$ given by (2.8) in the formula (2.2) of $u(y)$ with $y > 0$ one gets

$$\begin{aligned} u(y) &= \frac{y^{2\sigma}}{4^\sigma \Gamma(\sigma)} \int_0^\infty \frac{e^{-y^2/(4t)}}{t^{1+\sigma}} \int_0^\infty W^\alpha \left(\frac{e^{-s^2/(4t)}}{\sqrt{\pi t}} \right) \mathcal{C}_\alpha(s) f \, ds \, dt, \\ &= \frac{y^{2\sigma}}{4^\sigma \sqrt{\pi}\Gamma(\sigma)} \int_0^\infty W^\alpha \left(\int_0^\infty \frac{e^{-(y^2+s^2)/(4t)}}{t^{\sigma+1/2}} \frac{dt}{t} \right) \mathcal{C}_\alpha(s) f \, ds \\ &= \frac{y^{2\sigma}}{4^\sigma \sqrt{\pi}\Gamma(\sigma)} \int_0^\infty W^\alpha \left(\int_0^\infty e^{-r} \left(\frac{4r}{y^2+s^2} \right)^{\sigma+1/2} \frac{dr}{r} \right) \mathcal{C}_\alpha(s) f \, ds \\ &= \frac{2\Gamma(\sigma+1/2)}{\sqrt{\pi}\Gamma(\sigma)} \int_0^\infty W^\alpha \left(\frac{y^{2\sigma}}{(y^2+s^2)^{\sigma+1/2}} \right) \mathcal{C}_\alpha(s) f \, ds, \end{aligned}$$

as it was wanted.

Let us now assume that $f \in \mathcal{D}(A)$. Putting (2.8) in (2.5) we have for every $y > 0$ that

$$u(y) - f = \mu_\sigma \int_0^\infty W^\alpha \left(\int_0^\infty \frac{e^{-s^2/(4t)}(e^{-y^2/(4t)} - 1)}{t^{1/2-\sigma}} \frac{dt}{t} \right) \mathcal{C}_\alpha(s)(-A)^\sigma f \, ds,$$

where $\mu_\sigma = (\sqrt{\pi}\Gamma(\sigma))^{-1}$.

In order to compute $F(s,y) := \displaystyle\int_0^\infty e^{-s^2/(4t)}(e^{-y^2/(4t)}-1)t^{\sigma-3/2}dt$, note that this integral is analytic in $\Re\sigma < 3/2$. For a while, suppose that σ is such that $0 < \sigma < 1/2$. Then, for $s, y > 0$,

$$\begin{aligned} F(s,y) &= \int_0^\infty \frac{e^{-(s^2+y^2)/(4t)}}{t^{1/2-\sigma}} \frac{dt}{t} - \int_0^\infty \frac{e^{-s^2/(4t)}}{t^{1/2-\sigma}} \frac{dt}{t} \\ &= 4^{(1/2)-\sigma}\Gamma(1/2-\sigma) \left((s^2+y^2)^{\sigma-1/2} - (s^2)^{\sigma-1/2} \right). \end{aligned}$$

If $\sigma = 1/2$, $a = (s^2+y^2)/4$ and $b = s^2/4$ we get by partial derivation in a and b under the integral that

$$\begin{aligned} F(s,y) &= \int_0^\infty (e^{-a/t} - e^{-b/t}) \frac{dt}{t} = \int_0^\infty (e^{-ar} - e^{-br}) \frac{dr}{r} \\ &= \log(b/a) = \log\left(\frac{s^2}{y^2+s^2} \right), \quad \text{for all } s, y > 0. \end{aligned}$$

This proves the theorem. □

2.2.3. *Boundary values as integrated groups.* Boundary values of analytic semigroups in $\mathbb{C}^+$ give rise to important integrated groups or semigroups. The following theorem is [11, Corollary 5.3], see also [12, Theorem 2.3].

Theorem 2.9. *Let $\nu \geq 0$ and let H be the infinitesimal generator of an analytic semigroup $T(z)$ in $\mathbb{C}^+$. The following assertions are equivalent.*

(a) For all $\alpha > \nu$ there exists $\tau \geq 0$ such that the semigroup $T(z)$ satisfies estimate (2.6) with some constant $C_{\alpha,\tau}$.

(b) For all $\alpha > \nu$ the operator iH generates a globally tempered α-times integrated group $T_\alpha(it)$, $t \in \mathbb{R}$, of growth α.

According to this result, Theorem 1.1 applies to $A = -iL$ for L any of the operators included in Examples 2.6. These cases deserve a particular digression, which is done in the next subsection.

2.3. **Operators iH.** Let H be the infinitesimal generator of an analytic semigroup in $\mathbb{C}^+$ satisfying the estimate (2.6) and let $\alpha > \nu$. Then, by Theorem 2.9, iH generates an α-times integrated group $T_\alpha(it)$ such that

$$\|T_\alpha(it)\| \le C|t|^\alpha, \quad t \in \mathbb{R}.$$

Let us consider the extension problem (1.5) for $A = iH$ and $\sigma = 1/2$: $u_{yy} = -iHu$, $u(0) = f$. It is very simple to find a solution of this problem. Actually, it is enough to take the solution $v(z) = e^{-z\sqrt{-H}}f$ ($z \in \mathbb{C}^+$) to the problem (1.5) for $A = \sqrt{-H}$, $\sigma = 1/2$, and make $u(y) := v((\sqrt{2}/2)(1+i)y)$, $y \in \mathbb{R}$. Clearly, the existence of this solution is possible because the semigroup $e^{-t\sqrt{-H}}$, $t > 0$, admits an analytic extension to $\mathbb{C}^+$.

Let now assume that $\sigma \neq 1/2$. Once again the function $u(y) := v((\sqrt{2}/2)(1+i)y)$ is the *formal* solution in Theorem 1.1 of (1.5) for the integrated group $T_\alpha(iy)$ in this case, if v is the solution in Theorem 2.2 of (1.1) for the semigroup $e^{-t\sqrt{-H}}$. However, to give a sense to the function above one needs that $e^{-t\sqrt{-H}}$ admits an extension up to the boundary of $S_{\pi/4}$ at least, and that $e^{-t\sqrt{-H}}$ is bounded on $(0,\infty)$ (for which τ must be zero in (2.6)). Thus we get the following corollary.

Corollary 2.10. *Let H be the generator of an analytic semigroup e^{zH} satisfying, for some $\alpha, \tau \ge 0$,*

$$\|e^{zH}\| \le Ce^{\tau \Re z}\left(\frac{|z|}{\Re z}\right)^\alpha, \quad \forall z \in \mathbb{C}^+.$$

Let $T_\alpha(it)$ be the globally tempered α-times integrated group generated by iH. Let $\sigma \neq 1/2$. Then

(i) *the problem*

$$\left\{ \begin{array}{l} u_{yy} + \frac{1-2\sigma}{y} u_y = -iHu; \\ u(0) = f \quad f \in X; \end{array} \right. \tag{2.9}$$

has a solution given by formula (1.6), and also by (1.8) or (1.9) if $f \in \mathcal{D}(H)$, with $T_\alpha(it)$, $t \in \mathbb{R}$, in the integrals;

(ii) *in the case when $\tau = 0$ and $f \in \mathcal{D}(\sqrt{-H})$ that solution u coincides with*

$$u(y) = v((\sqrt{2}/2)(1+i)y), \quad y \in \mathbb{R},$$

where v is the solution to the problem

$$\left\{ \begin{array}{l} v_{yy} + \frac{1-2\sigma}{y}\, v_y = \sqrt{-H}v; \\ u(0) = f \quad f \in \mathcal{D}(\sqrt{-H}), \end{array} \right. \tag{2.10}$$

given by formulae (2.4) or (2.5) applied to the semigroup $e^{-z\sqrt{-H}}$.

The case $\sigma = 1/2$ is trivial, as seen prior to the theorem.

Remark 2.11. Since an operator H as in the corollary above generates a globally tempered cosine family $\mathcal{C}_\beta(s)$, the solution u to equation (2.9) can be also expressed as $u(y) = v((\sqrt{2}/2)(1+i)y)$, for $y \in \mathbb{R}$, where v is the solution of (2.10) given by the integral formulas (1.10), for any $f \in X$, and (1.11) associated to $\mathcal{C}_\beta(s)$.

Thus the corollary applies to all the examples $H = -L$ listed in Example 2.6. Among them we have the Schrödinger operator $i\Delta$ which is related to the Schrödinger equation of quantum mechanics [22]. In this case one can obtain solutions to the degenerate equation with complex coefficients

$$\left\{ \begin{array}{ll} i\Delta u + \frac{1-2\sigma}{y} u_y + u_{yy} = 0 = \operatorname{div}(M \cdot \nabla u), & \text{in } \mathbb{R}^n \times (0,\infty); \\ u(x,0) = f(x), & \text{on } \mathbb{R}^n; \end{array} \right.$$

with the $(n+1)\times(n+1)$ diagonal matrix $M = \operatorname{diag}(i,\ldots,i,y^{1-2\sigma})$, which, by Theorem 1.1, satisfy the L^p estimate

$$\sup_{y>0} \|u(y)\|_{L^p(\mathbb{R}^n)} \le C\,\|f\|_{L^p(\mathbb{R}^n)}\,.$$

Note that the coefficients of the equation above are not bounded, so the theory of elliptic equations with complex bounded measurable coefficients does not apply to it.

Fractional Schrödinger equations of the form $v_t = (-i\Delta)^\sigma v$, $v(x,0) = f(x)$, can be studied as limit cases when $z \to i$ of equations $v_t = (-z\Delta)^\sigma v$ for $\Re z > 0$, see [22, p. 283]. By Theorem 1.1 such a fractional Schrödinger equation can be written in the following equivalent form for a function $u = u(x,t,y)$:

$$\begin{cases} i\Delta u + \frac{1-2\sigma}{y} u_y + u_{yy} = 0, & \text{in } \mathbb{R}^n \times \mathbb{R} \times (0,\infty); \\ u_t(x,t,0) + \frac{1}{2\sigma c_\sigma} \lim\limits_{y\to 0^+} y^{1-2\sigma} u_y(x,t,y) = 0, & \text{in } \mathbb{R}^n \times \mathbb{R}. \end{cases}$$

Note that the operators H considered in the preceding lines have all real spectrum. In other words, they are particular cases of operators with pure imaginary symbols. We discuss this more general class of operators in the next subsection.

2.4. **Operators with pure imaginary symbol.** Conditions for operators A with purely imaginary symbols to generate integrated semigroups have been thoroughly studied by M. Hieber in a series of papers; see [17, 19, 18, 21, 20], or [1, Part C]. As a consequence of such an investigation we can apply our Theorem 1.1 to the following classes.

(1) Operators A for which their symbols $\widehat{A}$ are given by $\widehat{A}(\xi) = iq(\xi)$, $\xi \in \mathbb{R}^N$, where q is a real elliptic homogeneous polynomial or a real homogeneous C^∞ function on $\mathbb{R}^N \setminus \{0\}$ such that $\xi = 0$ if $q(\xi) = 0$. They generate globally tempered integrated semigroups; see [17, Theorem 4.2].

This result provides us with a lot of examples. In particular we can emphasize on the linear operator $\partial_x^3 = \partial^3/\partial x^3$ associated to the general Korteweg–de Vries equation,

$$\partial_t u = -\partial_x^3 u + \partial_x P(u), \quad (x,t) \in \mathbb{R}^2, \tag{2.11}$$

where $P(u)$ is some polynomial ($P(u) = 6u$ for the usual Korteweg–de Vries equation). It is known that solutions of equation (2.11) can be constructed out from solutions of the equation

$$\partial_t u = -\partial_x^3 u, \quad (x,t) \in \mathbb{R}^2, \tag{2.12}$$

see [33] and references therein. (In turn, a solution to the linearized KdV equation $v_t = -v_{xxx}$ is given by $v(x,t) = t^{-1/3} w(x/(3t)^{1/3})$ where w is a solution of the homogeneous Airy equation $w_{xx} - xw = 0$ [34, pp. 129, 130]).

By the aforementioned results of Hieber we have that the operator ∂_x^3 is the generator of a globally tempered α-times integrated semigroup on $L^p(\mathbb{R})$ for $\alpha > |1/p - 1/2|$ [17, p. 15]. Hence Theorem 1.1 holds also true for that operator, so that ∂_x^3 can be added to our examples.

Also in this case, Theorem 1.1 implies the following corollary.

Corollary 2.12. *The problem*

$$\begin{cases} \frac{1-2\sigma}{y} u_y + u_{yy} = u_{xxx}, & \textit{in } \mathbb{R} \times \mathbb{R} \times (0,\infty); \\ u_t(x,t,0) - \dfrac{1}{2\sigma c_\sigma} \lim\limits_{y\to 0^+} y^{1-2\sigma} u_y(x,t,y) = 0, & \textit{in } \mathbb{R} \times \mathbb{R}; \end{cases}$$

is equivalent to the fractional Korteweg–de Vries equation

$$\begin{cases} w_t = (-\partial_{xxx})^\sigma w, & \textit{in } \mathbb{R}^2,\ 0 < \sigma < 1; \\ w(x,0) = f(x), & \textit{on } \mathbb{R}. \end{cases}$$

with $w(x,t) \equiv u(x,t,y)$.

(2) Operators $A = P(D)$ on $L^p(\mathbb{R}^N)^m$, with $1 < p < \infty$, arising in symmetric hyperbolic systems like Maxwell‘s equations, elastic waves in homogeneous medium, or neutrino equation. If the symbol of these operators is homogeneous then $P(D)$ generates a globally tempered integrated semigroup. See [18, Corollary 3.2 and Section 4].

(3) Operators with symbols which are homogeneous coercive polynomials on $\mathbb{R}^N$ and have spectrum contained in a half-plane [21, Theorem 5.7].

(4) Pseudodifferential operators on L^p with spectrum within a half-plane and homogeneous symbol [20, Theorem 4.6 and Remark 4.7 b)].

3. Functions in convolution Sobolev algebras

For $\alpha > 0$ we define the Banach space of Sobolev type $\mathcal{T}^{(\alpha)}(t^\alpha)$ as the completion of the test function space $C_c^\infty[0,\infty)$ in the norm

$$\|\varphi\|_{(\alpha)} := \frac{1}{\Gamma(\alpha+1)} \int_0^\infty |W^\alpha \varphi(t)| t^\alpha \, dt, \quad \varphi \in C_c^\infty[0,\infty).$$

Here $W^\alpha \varphi$ is the Weyl fractional derivative of φ of order α given by the formulae

$$W^{-\beta}\varphi(s) := \frac{1}{\Gamma(\beta)} \int_s^\infty (t-s)^{\beta-1} \varphi(t)\, dt, \quad s \geq 0,$$

and

$$W^\alpha \varphi(s) := (-1)^n \frac{d^n}{ds^n} W^{-\beta}\varphi(s), \quad s \geq 0,$$

where $\beta = n - \alpha$, with $n = [\alpha] + 1$. In fact, the mapping from $C_c^\infty[0,\infty)$ into itself given by $\varphi(t) \longmapsto t^\alpha W^\alpha \varphi(t)$ extends to a Banach isomorphism from $\mathcal{T}^{(\alpha)}(t^\alpha)$ into $L^1(\mathbb{R}^+)$, so that a function φ belongs to $\mathcal{T}^{(\alpha)}(t^\alpha)$ if and only if there exists $g \in L^1(\mathbb{R}^+, t^\alpha\, dt)$ such that

$$\varphi(s) = \frac{1}{\Gamma(\alpha)} \int_s^\infty (t-s)^{\alpha-1} g(t)\, dt, \quad s > 0.$$

Then we put $g := W^\alpha \varphi$. Note that $W^n = (-1)^n (d^n/dt^n)$ for every $n \in \mathbb{N} \cup \{0\}$ and that $W^{\alpha+\beta}\varphi = W^\alpha(W^\beta \varphi)$ whenever the composition has sense.

We also have the inclusions $C_c^\infty[0,\infty) \hookrightarrow \mathcal{S}[0,\infty) \hookrightarrow \mathcal{T}^{(\beta)}(t^\beta) \hookrightarrow \mathcal{T}^{(\alpha)}(t^\alpha) \hookrightarrow \mathcal{T}^{(0)}(t^0) \equiv L^1(\mathbb{R}^+)$, whenever $\beta > \alpha > 0$, where $\mathcal{S}[0,\infty)$ is the restriction of the Schwartz's class $\mathcal{S}(\mathbb{R})$ to $[0,\infty)$. These inclusions are continuous for the usual topologies in $C_c^\infty[0,\infty)$ and $\mathcal{S}[0,\infty)$, and the norms $\|\cdot\|_{(\alpha)}$, $\alpha \geq 0$.

Furthermore, $\mathcal{T}^{(\alpha)}(t^\alpha)$ is a Banach algebra with product defined by the convolution on $\mathbb{R}^+$, which in this paper means that the convolution is jointly continuous with respect to the norm $\|\cdot\|_{(\alpha)}$. See [13, 14, 26] for these and some other properties of the Banach algebras $\mathcal{T}^{(\alpha)}(t^\alpha)$. We will refer to these algebras here as (convolution) Sobolev algebras.

In the remainder of this section we give some properties of three functions $b^{\sigma,z}$, $B^{\sigma,z}$ and h^σ, in connection with the Sobolev algebras. Such functions are important in the proof of Theorem 1.1, see Section 5 below.

Given $\sigma > 0$, define

$$b^{\sigma,z}(t) := \frac{z^{2\sigma}}{4^\sigma \Gamma(\sigma)} \frac{e^{-z^2/(4t)}}{t^{1+\sigma}}, \quad t > 0,\ z \in S_{\pi/4}. \tag{3.1}$$

Obviously, $b^{\sigma,z} \in L^1(\mathbb{R}^+)$ for all $\sigma > 0$, $z \in S_{\pi/4}$. Notice that, for $\sigma = 1/2$, the family $b^{1/2,z}$ is the well known backwards heat semigroup (see [36, p. 262], for instance, for a proof of the semigroup property). Next we point out several identities concerning derivatives of the function $b^{\sigma,z}$. In particular we will see that $b^{\sigma,z}$ belongs to the Sobolev algebras $\mathcal{T}^{(\alpha)}(t^\alpha)$ for any order of derivation α.

Lemma 3.1. *Let $z \in S_{\pi/4}$, $\sigma > 0$ and $n \in \mathbb{N}$.*

(i) $\displaystyle \frac{\partial^n}{\partial t^n} b^{\sigma,z}(t) = \left(\sum_{j=0}^n d_{j,n}^\sigma \frac{z^{2j}}{4^j t^{j+n}} \right) b^{\sigma,z}(t)$, *with $d_{j,n}^\sigma \in \mathbb{R}$. More precisely, for $n = 1$,*

$$\frac{\partial}{\partial t} b^{\sigma,z}(t) = \left(\frac{z^2}{4t^2} - \frac{1+\sigma}{t} \right) b^{\sigma,z}(t).$$

(ii) $\displaystyle \frac{\partial^n}{\partial z^n} b^{\sigma,z}(t) = \left(\sum_{j=0}^n c_{j,n}^\sigma \frac{z^{2j-n}}{t^j} \right) b^{\sigma,z}(t)$, *with $c_{j,n}^\sigma \in \mathbb{R}$. More precisely, for $n = 1, 2$,*

$$\frac{\partial}{\partial z} b^{\sigma,z}(t) = \left(\frac{2\sigma}{z} - \frac{z}{2t} \right) b^{\sigma,z}(t)$$

and

$$\frac{\partial^2}{\partial z^2}\, b^{\sigma,z}(t) = \left(\frac{2\sigma(2\sigma-1)}{z^2} - \frac{4\sigma+1}{2t} + \frac{z^2}{4t^2}\right) b^{\sigma,z}(t).$$

(iii) *The function* $b^{\sigma,z}$ *satisfies the differential equations*

$$2t\frac{\partial}{\partial t}\, u(z,t) + z\frac{\partial}{\partial z}\, u(z,t) = -2u(z,t),$$

and

$$\frac{\partial^2}{\partial z^2}\, u(z,t) + \frac{1-2\sigma}{z}\,\frac{\partial}{\partial z}\, u(z,t) = \frac{\partial}{\partial t}\, u(z,t),$$

with the boundary conditions $\lim_{t\to 0^+} u(z,t) = \lim_{t\to\infty} u(z,t) = 0$.

Proof. We just compute $\dfrac{\partial^n}{\partial t^n}\, b^{\sigma,z}$. The remainder of the proof is left to the interested reader. Put $a = -z^2/4$. For every $m \in \mathbb{N}$ and $p = 1, \ldots, m$ there are constants $c_{p,m} \in \mathbb{R}$, such that

$$(e^{at^{-1}})^{(m)}(t) = (-1)^m \sum_{p=1}^{m} c_{p,m}(a/t)^p t^{-m} e^{at^{-1}} = p_m(a/t)t^{-m}e^{at^{-1}}, \tag{3.2}$$

where p_m is a polynomial of degree m, with $p_m(0) = 0$. Hence, for all $n \in \mathbb{N}$ and $\sigma > 0$,

$$(e^{at^{-1}} t^{-(\sigma+1)})^{(n)}(t) = (-1)^n \sum_{m=0}^{n} c_{m,n,\sigma}\, p_m(a/t) t^{-(\sigma+n+1)} e^{at^{-1}}, \tag{3.3}$$

with $p_0 \equiv 1$, where $c_{m,n,\sigma} = \binom{n}{m}\binom{\sigma+n-m}{n-m}(n-m)!$. By reordering terms, we get

$$(e^{at^{-1}} t^{-(\sigma+1)})^{(n)}(t) = \sum_{k=0}^{n} d^{\sigma}_{k,n}(a/t)^k t^{-(\sigma+n+1)} e^{at^{-1}}. \tag{3.4}$$

Therefore, $\dfrac{\partial^n}{\partial t^n}\, b^{\sigma,z}(t) = \left(\displaystyle\sum_{j=0}^{n} d^{\sigma}_{j,n}\frac{z^{2j}}{4^j t^{j+n}}\right) b^{\sigma,z}(t)$, for some real constants $d^{\sigma}_{j,n}$, as claimed in the statement. □

Put $e_\varepsilon(t) := e^{-\varepsilon t}$ for all $\varepsilon, t > 0$.

Proposition 3.2. *Let* $(b^{\sigma,z})_{z\in S_{\pi/4}}$ *be the family of functions defined in* (3.1).

(i) *For every* $N \in \mathbb{N}$, *we have* $(b^{\sigma,z})_{z\in S_{\pi/4}} \subseteq \mathcal{T}^{(N)}(t^N)$ *with*

$$\|b^{\sigma,z}\|_{(N)} \le C_{N,\sigma}\left(\frac{|z|^2}{\Re(z^2)}\right)^{N+\sigma}, \quad z \in S_{\pi/4}.$$

Moreover, the map $z \longmapsto b^{\sigma,z}$, $S_{\pi/4} \to \mathcal{T}^{(N)}(t^N)$ *is analytic. In consequence,* $(b^{\sigma,z})_{z\in S_{\pi/4}}$ *is an analytic family of functions in* $\mathcal{T}^{(\alpha)}(t^\alpha)$, *for all* $\alpha > 0$.

(ii) *For every* $\alpha > 0$, *the equality*

$$\frac{\partial^2}{\partial z^2}\, b^{\sigma,z} + \frac{1-2\sigma}{z}\,\frac{\partial}{\partial z}\, b^{\sigma,z} = \frac{\partial}{\partial t}\, b^{\sigma,z}, \quad z \in S_{\pi/4},$$

holds in the Banach algebra $\mathcal{T}^{(\alpha)}(t^\alpha)$.

(iii) *For every* $\varepsilon > 0$ *and* $\alpha > 0$ *we have* $b^{\sigma,z}e_\varepsilon \in \mathcal{T}^{(\alpha)}(t^\alpha)$ *and, if* $0 < \sigma < 1$ *then* $\lim_{\varepsilon\to 0^+} b^{\sigma,z}e_\varepsilon = b^{\sigma}_z$ *in the norm of* $\mathcal{T}^{(\alpha)}(t^\alpha)$, *and uniformly on compact subsets of* $S_{\pi/4}$.

Proof. (i) Take $z \in S_{\pi/4}$, $\sigma > 0$ and $N \in \mathbb{N}$. By Lemma 3.1*(i)*,

$$\begin{aligned}\|b^{\sigma,z}\|_{(N)} &\le \sum_{j=0}^{N} |d^{\sigma}_{j,N}| 4^{-(j+\sigma)} \frac{|z|^{2(\sigma+j)}}{N!\Gamma(\sigma)} \int_0^\infty t^{-(1+j+\sigma)} |e^{-z^2/(4t)}|\, dt \\ &= \sum_{j=0}^{N} \frac{2|d^{\sigma}_{j,N}|\, |z|^{2(\sigma+j)}}{N!\Gamma(\sigma)\Re(z^2)^{\sigma+j}} \int_0^\infty u^{2(\sigma+j)-1} e^{-u^2} du \\ &\le C_{N,\sigma} \left(\frac{|z|^2}{\Re(z^2)}\right)^{N+\sigma},\end{aligned}$$

where we have used the change of variable $u^2 = \Re(z^2)/(4t)$ in the equality. Now we apply that $\mathcal{T}^{(N)}(t^N) \hookrightarrow \mathcal{T}^{(\alpha)}(t^\alpha)$ if $N \ge \alpha$ to conclude that $b^{\sigma,z} \in \mathcal{T}^{(\alpha)}(t^\alpha)$.

The analyticity of the map $z \longmapsto b^{\sigma,z}$, $z \in S_{\pi/4} \to \mathcal{T}^{(\alpha)}(t^\alpha)$, follows from its continuity (use the dominated convergence theorem) and Morera's theorem.

(ii) The assertion holds because, for $z \in S_{\pi/4}$, the functions $\dfrac{\partial^2}{\partial z^2} b^\sigma_z$, $\dfrac{\partial}{\partial z} b^\sigma_z$ and $\dfrac{\partial}{\partial t} b^\sigma_z$ belong to $\mathcal{T}^{(\alpha)}(t^\alpha)$ for every $\alpha \ge 0$ (see Lemma 3.1).

(iii) Write $a = -z^2/4$. It follows by (3.3) that, for $n = 0, 1, \ldots, N$ and $t > 0$,

$$\begin{aligned}|(e^{at^{-1}} t^{-(\sigma+1)})^{(n)}(t)| &\le \sum_{k=0}^{n} |d^{\sigma}_{k,n}| |a|^k t^{-k} e^{(\Re a/2)t^{-1}} t^{-(\sigma+n+1)} e^{(\Re a/2)t^{-1}} \\ &\le \sum_{k=0}^{n} |d^{\sigma}_{k,n}| |a|^k \left(\frac{2k}{|\Re a|}\right)^k e^{-k} t^{-(\sigma+n+1)} e^{(\Re a/2)t^{-1}}.\end{aligned}$$

Hence, if K is a compact subset of $S_{\pi/4}$ then there exist constants $C_{\sigma,K}, M_K > 0$ such that

$$|(b^{\sigma,z})^{(n)}(t)| \le C_{\sigma,K} t^{-(\sigma+n+1)} e^{-M_K t^{-1}}, \quad t > 0,\ z \in K.$$

Then, if $0 < \sigma < 1$, for every $z \in K$ we have

$$\begin{aligned}\|b^{\sigma,z} e_\varepsilon - b^{\sigma,z}\|_{(N)} &\le \sum_{n=0}^{N-1} \int_0^\infty |(b^{\sigma,z})^{(n)}(t)| \varepsilon^{N-n} e^{-\varepsilon t} t^N dt \\ &\quad + \sum_{n=0}^{N-1} \int_0^\infty |(b^{\sigma,z})^{(N)}(t)| (1 - e^{-\varepsilon t}) t^N dt \\ &\le C_{\sigma,K} \left[\sum_{n=0}^{N-1} \varepsilon^{N-n} \int_0^\infty t^{N-n} e^{-\varepsilon t} \frac{dt}{t^{1+\sigma}} + \int_0^\infty e^{-M_K/t} \left(1 - e^{-\varepsilon t}\right) \frac{dt}{t^{\sigma+1}} \right] \\ &= C_{\sigma,K} \left[\sum_{n=0}^{N-1} \Gamma(N-\sigma-n) \varepsilon^\sigma + \int_0^\infty e^{-M_K/t} \left(1 - e^{-\varepsilon t}\right) \frac{dt}{t^{\sigma+1}} \right].\end{aligned}$$

From here, by dominated convergence in the last integral we obtain that $\sup_{z \in K} \|b^{\sigma,z} e_\varepsilon - b^{\sigma,z}\|_{(N)} \to 0$, as $\varepsilon \to 0^+$. □

Remark 3.3. The Laplace transform of $b^{\sigma,z}$ can be expressed as

$$\mathcal{L}(b^{\sigma,z})(w) = \frac{2^{1-\sigma}}{\Gamma(\sigma)} (z\sqrt{w})^\sigma K_\sigma\left(z\sqrt{w}\right), \quad z \in S_{\pi/4},\ w \in \mathbb{C}^+, \tag{3.5}$$

where K_σ is the modified Bessel function of the third kind, see [24, p. 119]. Then the function K_σ is implicitly subject to the following estimates: for every $\alpha > 0$, $b^{\sigma,z}$ belongs to $\mathcal{T}^{(\alpha)}(t^\alpha)$ by Proposition

3.2, and therefore its Laplace transform lies in the algebra $\mathcal{A}_0^{(\alpha)}(\mathbb{C}^+)$ introduced in [14]. This means that $\mathcal{L}(b^{\sigma,z})$ is continuous in $\overline{\mathbb{C}}^+$, holomorphic in $\mathbb{C}^+$, and it satisfies

$$\lim_{w\to\infty} w^\beta \mathcal{L}(b^{\sigma,z})^{(\beta)}(w) = 0, \quad 0 \le \beta \le \alpha;$$

$$\lim_{w\to 0} w^\beta \mathcal{L}(b^{\sigma,z})^{(\beta)}(w) = 0, \quad 0 < \beta \le \alpha.$$

The space $\mathcal{A}_0^{(\alpha)}(\mathbb{C}^+)$ is a Banach algebra on $\mathbb{C}^+$ for the pointwise multiplication, which is formed by holomorphic functions possesing fractional derivatives up to the boundary of $\mathbb{C}^+$. For F in $\mathcal{A}_0^{(\alpha)}(\mathbb{C}^+)$, the notation $F^{(\beta)}$ refers to fractional derivative.

For $t, \sigma > 0$ and $z \in S_{\pi/4}$, define

$$B^{\sigma,z}(t) := 4^\sigma \left(\frac{t}{z}\right)^{2\sigma} b^{\sigma,z}(t) = \frac{1}{\Gamma(\sigma)} \frac{e^{-z^2/(4t)}}{t^{1-\sigma}}. \tag{3.6}$$

It is readily seen that

$$z^{1-2\sigma} \frac{\partial}{\partial z} B^{\sigma,z} = \frac{\sigma\Gamma(-\sigma)}{2^{2\sigma-1}\Gamma(\sigma)} b^{1-\sigma,z}, \quad z \in S_{\pi/4}, \tag{3.7}$$

where $0 < \sigma < 1$.

Besides the preceding relationships between the functions $b^{\sigma,z}$ and $B^{\sigma,z}$, we have the following identities involving $B^{\sigma,z}$, whose proof is left to prospective readers.

Lemma 3.4. *Let $z \in S_{\pi/4}$, $t > 0$ and $n \in \mathbb{N}$.*

(i) $\frac{\partial^n}{\partial t^n} B^{\sigma,z}(t) = \left(\sum_{j=0}^n k_{j,n}^\sigma \frac{z^{2j}}{4^j t^{j+n}}\right) B^{\sigma,z}(t)$, *with constants* $k_{j,n}^\sigma \in \mathbb{R}$ *and* $k_{0,n}^\sigma = (\sigma-1)\cdots(\sigma-n)$. *In particular,*

$$\frac{\partial}{\partial t} B^{\sigma,z}(t) = \left(\frac{z^2}{4t^2} - \frac{1-\sigma}{t}\right) B^{\sigma,z}(t).$$

(ii) $\frac{\partial}{\partial z} B^{\sigma,z}(t) = -\frac{z}{2t} B^{\sigma,z}(t)$, *and*

$$\frac{\partial^2}{\partial z^2} B^{\sigma,z}(t) = \left(\frac{z^2}{4t^2} - \frac{1}{2t}\right) B^{\sigma,z}(t).$$

(iii) *The function $B^{\sigma,z}$ satisfies the differential equations*

$$2t\frac{\partial}{\partial t} u(z,t) + z\frac{\partial}{\partial z} u(z,t) = -2(1-\sigma)u(z,t),$$

and

$$\frac{\partial^2}{\partial z^2} u(z,t) + \frac{1-2\sigma}{z} \frac{\partial}{\partial z} u(z,t) = \frac{\partial}{\partial t} u(z,t),$$

with the boundary conditions $\lim_{t\to 0^+} u(z,t) = \lim_{t\to\infty} u(z,t) = 0$.

Let h^σ denote the Heaviside-type function defined by

$$h^\sigma(t) := \frac{1}{\Gamma(\sigma)} t^{\sigma-1}, \quad t > 0.$$

Lemma 3.5. *For $\sigma > 0$ and $z \in S_{\pi/4}$, we have $B^{\sigma,z} = h^\sigma * b^{\sigma,z}$; namely,*

$$B^{\sigma,z}(s) = \frac{1}{\Gamma(\sigma)} \int_0^s (s-t)^{\sigma-1} b^{\sigma,z}(t)\, dt, \quad s > 0.$$

Proof. For positive σ, z and s we have

$$\frac{1}{4^\sigma\Gamma(\sigma)}\int_0^s (s-t)^{\sigma-1}e^{-z^2/(4t)}\,\frac{dt}{t^{1+\sigma}} = \frac{1}{4^\sigma\Gamma(\sigma)}\int_0^s \left(\frac{s}{t}-1\right)^{\sigma-1} e^{-z^2/(4t)}\,\frac{dt}{t^2}$$
$$= s^{\sigma-1}\,e^{-z^2/(4s)}\int_0^\infty r^\sigma e^{-z^2 r}\,\frac{dr}{r} = \frac{s^{\sigma-1}\,e^{-z^2/(4s)}}{z^{2\sigma}},$$

where we have done the change of variable $4r = t^{-1} - s^{-1}$ in the second equality. From that, one gets that $B^{\sigma,z} = h^\sigma * b^{\sigma,z}$ for $\sigma, z > 0$. Then this equality extends to $z \in S_{\pi/4}$ by the analytic continuation principle. ☐

Clearly, $B^{\sigma,z} \notin L^1(\mathbb{R}^+)$, which is not good for our needs. However, the additive perturbation of $B^{\sigma,z}$ implemented by h^σ fits very well in our setting. Recall that $e_\varepsilon(t) = e^{-\varepsilon t}$ for $\varepsilon, t > 0$.

Proposition 3.6. *Take* $0 < \sigma < 1$, $z \in S_{\pi/4}$.

(i) *The function* $B^{\sigma,z} - h^\sigma$, *which is given by*

$$(B^{\sigma,z} - h^\sigma)(t) = \frac{1}{\Gamma(\sigma)}\frac{e^{-z^2/(4t)} - 1}{t^{1-\sigma}}, \quad \textit{for } t > 0,$$

belongs to $\mathcal{T}^{(N)}(t^N)$ *for every* $N \in \mathbb{N}$, *and*

$$\|B^{\sigma,z} - h^\sigma\|_{(N)} \le C_{N,\sigma}\frac{|z|^{2N}}{(\Re(z^2))^{N-\sigma}}, \quad z \in S_{\pi/4}.$$

As a consequence, $(B^{\sigma,z} - h^\sigma)_{z\in S_{\pi/4}} \subseteq \mathcal{T}^{(\alpha)}(t^\alpha)$, *for every* $\alpha > 0$.

(ii) *For every* $\varepsilon > 0$ *and* $\alpha > 0$, *we have that* $(B^{\sigma,z} - h^\sigma)\,e_\varepsilon$ *is in* $\mathcal{T}^{(\alpha)}(t^\alpha)$, *and moreover*

$$\lim_{\varepsilon\to 0^+}(B^{\sigma,z} - h^\sigma)\,e_\varepsilon = (B^{\sigma,z} - h^\sigma)$$

in the norm of $\mathcal{T}^{(\alpha)}(t^\alpha)$.

Proof. (i) The expression of the function $B^{\sigma,z} - h^\sigma$ follows rightly from the definitions. As regards the estimate of the statement, it is readily obtained using Lemma 3.4 (i). Item (ii) follows as in the proof of Proposition 3.2. We skip the details. ☐

Remark 3.7. For any $w \in \mathbb{C}^+$, one has $\mathcal{L}(h^\sigma)(w) = w^{-\sigma}$. Hence, if one takes Laplace transforms in the convolution identity of Lemma 3.5, on account of (3.5), one obtains

$$\mathcal{L}(B^{\sigma,z})(w) = \frac{2^{1-\sigma}}{\Gamma(\sigma)}\left(\frac{z}{\sqrt{w}}\right)^\sigma K_\sigma\left(z\sqrt{w}\right), \quad w \in \mathbb{C}^+.$$

Since $B^{\sigma,z} - h^\sigma$ lies in $\mathcal{T}^{(\alpha)}(t^\alpha)$, we have that the function

$$\mathcal{L}(B^{\sigma,z} - h^\sigma)(w) = \left(\frac{2^{1-\sigma}}{\Gamma(\sigma)}(z\sqrt{w})^\sigma K_\sigma\left(z\sqrt{w}\right) - 1\right) w^{-\sigma}, \quad w \in \mathbb{C}^+,$$

is subject for all $\alpha > 0$ to the same type of estimates as those mentioned in Remark 3.3.

In order not to make this section too long we stop here giving the primary properties of significant elements of the Banach algebra $\mathcal{T}^{(\alpha)}(t^\alpha)$. Properties of other functions in $\mathcal{T}^{(\alpha)}(t^\alpha)$ which are directly related with specific operations on operators on Banach spaces are given in the next section.

4. Fractional powers and perturbations of generators of α-times integrated semigroups

Let $\alpha > 0$. The notion of α-times integrated semigroup was introduced by M. Hieber in [19], see also [1, Section 3.17]. Here we consider α-times integrated semigroups $(T_\alpha(t))_{t\ge 0}$ which are globally tempered. Let us recall the corresponding definition given in the Introduction, which is implemented by formula (1.3) relating an α-times integrated semigroup with its generator:

$$(\lambda - A)^{-1}f = \lambda^\alpha\int_0^\infty e^{-\lambda t}T_\alpha(t)f\,dt, \quad \Re\lambda > 0,\ f \in X. \tag{4.1}$$

Each uniformly bounded C_0 semigroup $(T(t))_{t\geq 0}$ is a tempered α-times integrated semigroup for $\alpha = 0$. Other (non-trivial) examples of integrated semigroups have been given in Section 2. We will make usage of the following formulas or properties of integrated semigroups.

- For $f \in X$ and $\beta > \alpha$ define

$$T_\beta(t)f := \frac{1}{\Gamma(\beta-\alpha)} \int_0^t (t-s)^{\beta-\alpha-1} T_\alpha(s)f\,ds. \tag{4.2}$$

Then $T_\beta(t)$ is a globally tempered β-times integrated semigroup in $\mathcal{B}(X)$ with generator A as well.

- For $f \in \mathcal{D}(A)$,

$$T_\alpha(t)f - \frac{t^\alpha}{\Gamma(\alpha+1)} f = \int_0^t T_\alpha(s)Af\,ds = T_{\alpha+1}(t)Af, \quad t \geq 0; \tag{4.3}$$

see [19, Proposition 2.4].

- Clearly from the definition, the generator A of a globally tempered α-times semigroup satisfies

$$\left\|\lambda(\lambda - A)^{-1}\right\| \leq C, \quad \text{for all } \lambda > 0, \tag{4.4}$$

and then, for $0 < \sigma < 1$, the fractional power $(-A)^\sigma$ (in the Balakrishnan sense) exists and is given by

$$(-A)^\sigma f = \frac{\sin(\pi\sigma)}{\pi} \int_0^\infty \lambda^{\sigma-1}(\lambda - A)^{-1}(-Af)\,d\lambda, \quad f \in \mathcal{D}(A); \tag{4.5}$$

see [36, p. 260] or [25].

- For $\varepsilon > 0$ and $0 < \sigma < 1$, the operator $(\varepsilon - A)$ also satisfies the estimate (4.4) and then

$$\lim_{\varepsilon\to 0^+} (\varepsilon - A)^{-\sigma}(-A)^\sigma f = f, \quad f \in \mathcal{D}(A); \tag{4.6}$$

see [25, Theorem 2.1 and Theorem 3.2].

We now give an extension, for general globally tempered integrated semigroups, of a well-known formula for C_0-semigroups; see [36, p. 260].

Theorem 4.1. *Let A be the generator of a tempered α-times integrated semigroup $(T_\alpha(t))_{t\geq 0}$ and let σ be such that $0 < \sigma < 1$. Then, for $f \in \mathcal{D}(A)$,*

$$(-A)^\sigma f = \frac{\Gamma(\sigma+\alpha+1)}{\Gamma(-\sigma)\Gamma(1+\sigma)} \int_0^\infty \left(T_\alpha(t)f - \frac{t^\alpha}{\Gamma(\alpha+1)} f \right) \frac{dt}{t^{\sigma+\alpha+1}}.$$

Proof. Set $T_{\alpha+1}(t)f := \displaystyle\int_0^t T_\alpha(s)f\,ds$, $f \in X$, as above. Take $f \in \mathcal{D}(A)$. Since $T_{\alpha+1}(t)$ is globally tempered with exponent $\alpha + 1$, (4.1) applies so we have

$$(\lambda - A)^{-1} f = \lambda^{\alpha+1} \int_0^\infty e^{-\lambda t} T_{\alpha+1}(t)f\,dt, \quad \Re\lambda > 0. \tag{4.7}$$

Plugging (4.7) into (4.5) we see that the order of integration can be exchanged, since $\|T_{\alpha+1}(t)Af\| \leq C\,\|Af\|\,t^{\alpha+1}$, if $0 \leq t \leq 1$, and $\|T_{\alpha+1}(t)Af\| \leq C\,\|f\|\,t^\alpha$, if $t \geq 1$, by (4.3). Thus we obtain

$$\begin{aligned}
(-A)^\sigma f &= \frac{\sin(\pi\sigma)}{\pi} \int_0^\infty \lambda^{\sigma+\alpha} \int_0^\infty e^{-\lambda t} T_{\alpha+1}(t)(-Af)\,dt\,d\lambda \\
&= \frac{\sin(\pi\sigma)}{\pi} \int_0^\infty \lambda^{\sigma+\alpha} \int_0^\infty e^{-\lambda t} \left(\frac{t^\alpha}{\Gamma(\alpha+1)} f - T_\alpha(t)f \right) dt\,d\lambda \\
&= \frac{\Gamma(\sigma+\alpha+1)}{\Gamma(-\sigma)\Gamma(1+\sigma)} \int_0^\infty \left(T_\alpha(t)f - \frac{t^\alpha}{\Gamma(\alpha+1)} f \right) \frac{dt}{t^{\sigma+\alpha+1}}.
\end{aligned}$$

□

There is a useful connection between the algebras $\mathcal{T}^{(\alpha)}(t^\alpha)$, $\alpha \geq 0$, (introduced in the previous section) and globally temperated integrated semigroups $T_\alpha(t)$. The mapping $\pi_\alpha : \mathcal{T}^{(\alpha)}(t^\alpha) \to \mathcal{B}(X)$ defined by

$$\pi_\alpha(\varphi)f := \int_0^\infty W^\alpha \varphi(t)\, T_\alpha(t) f\, dt, \quad f \in X,\ \varphi \in \mathcal{T}^{(\alpha)}(t^\alpha), \tag{4.8}$$

is a bounded Banach algebra homomorphism. Also,

$$-A\pi_\alpha(\varphi)f = \pi_\alpha(\varphi')f + \varphi(0)f, \tag{4.9}$$

for every $\varphi \in C^1[0,\infty)$ such that $\varphi' \in \mathcal{T}^{(\alpha)}(t^\alpha)$. See [26, Theorem 3.1], where this is shown for φ in the Schwartz class. The proof works also in the case above.

We take advantage of the link between $T_\alpha(t)$ and π_α to establish a result about perturbations of α-times integrated semigroups. For $\varepsilon > 0$, it was proved by van Neerven and Straub that an ε-perturbation of an integrated semigroup of exponential growth is again an integrated semigroup of exponential growth, see [35, Proposition 3.3]. We show here that this property remains true if the exponential growth is replaced by polynomial, α-homogeneous, growth.

Theorem 4.2. *Let A be the generator of a tempered α-times integrated semigroup $T_\alpha(t) \subseteq \mathcal{B}(X)$. Take $\varepsilon > 0$.*

- (i) *The operator $A - \varepsilon$ is the generator of an α-times integrated semigroup $T_{\alpha,\varepsilon}(t)$ such that*
$$\|T_{\alpha,\varepsilon}(t)\| \leq C_\alpha t^\alpha, \quad t > 0,$$
where the constant C_α is independent of ε.
- (ii) *Moreover, for $t > 0$ and $f \in X$,*
$$T_{\alpha,\varepsilon}(t)f = e^{-\varepsilon t} T_\alpha(t)f + \sum_{k=1}^\infty \binom{\alpha}{k} \varepsilon^k \int_u^t \frac{(t-u)^{k-1}}{(k-1)!} e^{-\varepsilon u} T_\alpha(u) f\, du.$$

We give the proof of Theorem 4.2 divided into several steps. The first one consists of estimating a function $\phi^\alpha_{t,\varepsilon}$ (see (4.10) below) in $\mathcal{T}^{(\alpha)}(t^\alpha)$.

Lemma 4.3. *For $t > s \geq 0$,*

$$\frac{(t-s)^{\alpha-1}}{\Gamma(\alpha)} \left(e^{z(t-s)} - 1 \right) = \sum_{k=1}^\infty \binom{\alpha}{k} z^k \int_s^t \frac{(u-s)^{\alpha-1}(t-u)^{k-1}}{\Gamma(\alpha)(k-1)!} e^{z(t-u)} du.$$

Proof. By using the Taylor expansion of e^w it is clear that

$$\begin{aligned}
\frac{(t-s)^{\alpha-1}}{\Gamma(\alpha)} \left(e^{z(t-s)} - 1 \right) &= \sum_{j=0}^\infty \frac{z^{j+1}(t-s)^{\alpha+j}}{(j+1)!\Gamma(\alpha)} \\
&= \sum_{j=0}^\infty \frac{z^{j+1}}{\Gamma(\alpha) j!} \binom{\alpha+j}{j+1} \int_s^t (u-s)^{\alpha-1}(t-u)^j\, du.
\end{aligned}$$

On the other hand,

$$\begin{aligned}
&\sum_{k=1}^\infty \binom{\alpha}{k} z^k \int_s^t \frac{(u-s)^{\alpha-1}}{\Gamma(\alpha)} \frac{(t-u)^{k-1}}{(k-1)!} e^{z(t-u)}\, du \\
&\qquad = \sum_{k=0}^\infty \sum_{l=0}^\infty \binom{\alpha}{k+1} z^{k+l+1} \int_s^t \frac{(u-s)^{\alpha-1}}{\Gamma(\alpha)} \frac{(t-u)^{k+l}}{k!\, l!}\, du \\
&\qquad = \sum_{j=0}^\infty z^{j+1} \int_s^t \frac{(u-s)^{\alpha-1}}{\Gamma(\alpha)} \frac{(t-u)^j}{j!} \sum_{k=0}^j \binom{\alpha}{k+1} \binom{j}{k}\, du \\
&\qquad = \sum_{j=0}^\infty \frac{z^{j+1}}{\Gamma(\alpha) j!} \int_s^t (u-s)^{\alpha-1}(t-u)^j \binom{\alpha+j}{j+1}\, du,
\end{aligned}$$

where in the last equality we have applied the well-known Vandermonde identity (also known as Chu-Vandermonde identity). □

Proposition 4.4. *Take* $\alpha, \varepsilon > 0$ *and* $t \geq 0$. *Let* $\phi^{\alpha}_{t,\varepsilon}$ *be defined by*

$$\phi^{\alpha}_{t,\varepsilon}(s) := \sum_{k=1}^{\infty} \binom{\alpha}{k} \varepsilon^k \int_s^t \frac{(u-s)^{\alpha-1}}{\Gamma(\alpha)} \frac{(t-u)^{k-1}}{(k-1)!} e^{-\varepsilon u}\, du, \tag{4.10}$$

for $t \geq s \geq 0$, *and* $\phi^{\alpha}_{t,\varepsilon}(s) := 0$ *if* $s > t$. *Then* $\phi^{\alpha}_{t,\varepsilon} \in \mathcal{T}^{(\alpha)}(t^{\alpha})$ *with* $\|\phi^{\alpha}_{t,\varepsilon}\|_{(\alpha)} \leq t^{\alpha}/\Gamma(\alpha+1)$.

Proof. For $\alpha, \varepsilon > 0$, $t \geq 0$ and $0 \leq s \leq t$, write

$$F^{\alpha,k}_{t,\varepsilon}(s) := \int_s^t \frac{(u-s)^{\alpha-1}}{\Gamma(\alpha)} \frac{(t-u)^{k-1}}{(k-1)!} e^{-\varepsilon u}\, du,$$

so that $\phi^{\alpha}_{t,\varepsilon} = \sum_{k=1}^{\infty} \binom{\alpha}{k} \varepsilon^k F^{\alpha,k}_{t,\varepsilon}$. Then $W^{\alpha} F^{\alpha,k}_{t,\varepsilon}(s) = (t-s)_+^{k-1} e^{-\varepsilon s}/\Gamma(k)$, whence $F^{\alpha,k}_{t,\varepsilon} \in \mathcal{T}^{(\alpha)}(t^{\alpha})$ and $\|F^{\alpha,k}_{t,\varepsilon}\|_{(\alpha)} \leq t^{\alpha+k}/(k!\Gamma(\alpha))$.

Since the radius of convergence of the series $\displaystyle\sum_{k=1}^{\infty} \binom{\alpha}{k} \frac{z^k}{k!}$ is infinite, it follows that $\phi^{\alpha}_{t,\varepsilon} \in \mathcal{T}^{(\alpha)}(t^{\alpha})$. Moreover,

$$W^{\alpha}\phi^{\alpha}_{t,\varepsilon}(s) = \sum_{k=1}^{\infty} \binom{\alpha}{k} \varepsilon^k W^{\alpha} F^{\alpha}_{t,\varepsilon}(s) = \sum_{k=1}^{\infty} \binom{\alpha}{k} \varepsilon^k \frac{(t-s)_+^{k-1}}{(k-1)!} e^{-\varepsilon s}.$$

Thus $W^{\alpha}\phi^{\alpha}_{t,\varepsilon}(s) = 0$, for $s > t$, and $W^{\alpha}\phi^{\alpha}_{t,\varepsilon}(s) = (g_{\varepsilon,\alpha})'(t-s)e^{-\varepsilon s}$, for $0 < s < t$, where

$$g_{\varepsilon,\alpha}(u) = 1 + \sum_{k=1}^{\infty} \binom{\alpha}{k} \frac{(u\varepsilon)^k}{k!}, \quad u > 0.$$

By [35, Lemma 2.1], $g_{\varepsilon,\alpha}$ is an increasing function in $u > 0$, and therefore we have that $W^{\alpha}\phi^{\alpha}_{t,\varepsilon}(s) \geq 0$ if $0 \leq s < t$.

Hence

$$\begin{aligned}
\|\phi^{\alpha}_{t,\varepsilon}\|_{(\alpha)} &= \int_0^t \int_u^t \frac{(s-u)^{\alpha-1}}{\Gamma(\alpha)} W^{\alpha}\phi^{\alpha}_{t,\varepsilon}(s)\, ds\, du \\
&= \int_0^t \int_u^t \frac{(s-u)^{\alpha-1}}{\Gamma(\alpha)} \sum_{k=1}^{\infty} \binom{\alpha}{k} \varepsilon^k \frac{(t-s)^{k-1}}{(k-1)!} e^{-\varepsilon s}\, ds\, du \\
&= e^{-\varepsilon t} \int_0^t \sum_{k=1}^{\infty} \binom{\alpha}{k} \varepsilon^k \int_u^t \frac{(s-u)^{\alpha-1}}{\Gamma(\alpha)} \frac{(t-s)^{k-1}}{(k-1)!} e^{\varepsilon(t-s)}\, ds\, du \\
&= e^{-\varepsilon t} \int_0^t \frac{(t-u)^{\alpha-1}}{\Gamma(\alpha)} \left(e^{\varepsilon(t-u)} - 1\right) du \\
&= e^{-\varepsilon t} \int_0^t \frac{r^{\alpha-1}}{\Gamma(\alpha)} (e^{\varepsilon r} - 1)\, dr = e^{-\varepsilon t} \int_0^t \frac{r^{\alpha-1}}{\Gamma(\alpha)} \sum_{k=1}^{\infty} \frac{\varepsilon^k r^k}{k!} dr \\
&= \frac{e^{-\varepsilon t} t^{\alpha}}{\Gamma(\alpha)} \sum_{k=1}^{\infty} \frac{(\varepsilon t)^k}{k!(\alpha+k)} \leq \frac{t^{\alpha}}{\Gamma(\alpha+1)}
\end{aligned}$$

where we have used the identity $\displaystyle\frac{s^{\alpha}}{\Gamma(\alpha+1)} = \int_0^s \frac{(s-u)^{\alpha-1}}{\Gamma(\alpha)}\, du$ in the first equality, and Lemma 4.3 in the fourth equality. □

For $z \in \mathbb{C}$, $t \geq 0$ and $\alpha > 0$, define the function $R^{\alpha-1}_{t,z}$ by $R^{\alpha-1}_{0,z} = 0$, if $t = 0$, and

$$R^{\alpha-1}_{t,z}(s) = \frac{(t-s)^{\alpha-1}}{\Gamma(\alpha)} \chi_{(0,t)}(s) e^{-zs}, \quad s \geq 0,$$

for $t > 0$. Here χ_I denotes the indicator function of a given interval I.

Recall the definition of mutipliers of commutative Banach algebras as regards the algebra $\mathcal{T}^{(\alpha)}(t^\alpha)$. So, a multiplier of $\mathcal{T}^{(\alpha)}(t^\alpha)$ is a bounded linear mapping $\Lambda : \mathcal{T}^{(\alpha)}(t^\alpha) \to \mathcal{T}^{(\alpha)}(t^\alpha)$ such that $\Lambda(\varphi * g) = \varphi * \Lambda(g)$ for all $\varphi, g \in \mathcal{T}^{(\alpha)}(t^\alpha)$. It was shown in [13, Proposition 1.5] that the functions $R_t^{\alpha-1} := R_{t,0}^{\alpha-1}$, $t > 0$, acting by convolution, are multipliers of the Banach algebra $\mathcal{T}^{(\alpha)}(t^\alpha)$ such that

$$\|R_t^{\alpha-1}\| \leq C_\alpha t^\alpha, \quad t > 0.$$

We extend such a bound to the family of functions $(R_{t,\varepsilon}^{\alpha-1})_{t,\varepsilon>0}$, with a uniform bound independent of ε

Proposition 4.5. *For every $t, \varepsilon > 0$, the function $R_{t,\varepsilon}^{\alpha-1}$ defines by convolution a multiplier in $\mathcal{T}^{(\alpha)}(t^\alpha)$ such that, for some constant $C = C_\alpha$ independent of ε, $\|R_{t,\varepsilon}^{\alpha-1}\| \leq Ct^\alpha$, for all $t > 0$.*

Proof. By Lemma 4.3, $R_{t,\varepsilon}^{\alpha-1} = e^{-\varepsilon t} R_t^{\alpha-1} + \phi_{t,\varepsilon}^\alpha$, where $\phi_{t,\varepsilon}^\alpha \in \mathcal{T}^{(\alpha)}(t^\alpha)$ by Proposition 4.4. Also, $R_t^{\alpha-1}$ is a convolution multiplier of $\mathcal{T}^{(\alpha)}(t^\alpha)$ such that $\|R_t^{\alpha-1}\| \leq Ct^\alpha$ for every $t > 0$, see [13, Proposition 1.5]. Therefore $R_{t,\varepsilon}^{\alpha-1}$ defines by convolution a multiplier of $\mathcal{T}^{(\alpha)}(t^\alpha)$, and by Proposition 4.4, $\|R_{t,\varepsilon}^{\alpha-1}\| \leq e^{-\varepsilon t}\|R_t^{\alpha-1}\| + \|\phi_{t,\varepsilon}^\alpha\|_\alpha \leq C_\alpha t^\alpha$, as we wanted to show. ☐

Proof of Theorem 4.2. Let A be the generator of a globally tempered α-times integrated semigroup $T_\alpha(t)$. Then the bounded homomorphism $\pi_\alpha : \mathcal{T}^{(\alpha)}(t^\alpha) \to \mathcal{B}(X)$ satisfies $\pi_\alpha(\mathcal{T}^{(\alpha)}(t^\alpha))X = X$. The map π_α can be extended to the multiplier algebra of $\mathcal{T}^{(\alpha)}(t^\alpha)$ by defining

$$\pi_\alpha(\Lambda)f = \pi_\alpha(\Lambda(\varphi))g$$

for every multiplier Λ, where $\varphi \in \mathcal{T}^{(\alpha)}(t^\alpha)$ and $g \in X$ are such that $f = \pi_\alpha(\varphi)g$ (it is readily seen that the definition does not depend on the choice of φ and g). In such a way, one gets the definition of $\pi_\alpha(R_t^{\alpha-1})$ and, in fact,

$$\pi_\alpha(R_t^{\alpha-1}) = T_\alpha(t), \quad t > 0; \tag{4.11}$$

see [13].

By [35, Proposition 3.3], the perturbation A_ε is the generator of an α-times integrated semigroup $T_{\alpha,\varepsilon}(t)$ which is related with $T_\alpha(t)$ through the formula

$$T_{\alpha,\varepsilon}(t)f = e^{-\varepsilon t} T_\alpha(t) f + \sum_{k=1}^\infty \binom{\alpha}{k} \varepsilon^k \int_0^t \frac{(t-s)^{k-1}}{(k-1)!} e^{-\varepsilon s} T_\alpha(s) f \, ds$$

for every $t > 0$ and $f \in X$.

By Proposition 4.4 and (4.11), the formula above reads

$$T_{\alpha,\varepsilon}(t)f = e^{-\varepsilon t}\pi_\alpha(R_t^{\alpha-1})f + \pi_\alpha(\phi_{t,\varepsilon}^\alpha)f = \pi_\alpha(R_{t,\varepsilon}^{\alpha-1})f,$$

for every $t > 0$ and $f \in X$. Then by Proposition 4.5 and Proposition 4.4 we conclude that

$$\|T_{\alpha,\varepsilon}(t)\| \leq C_\alpha t^\alpha, \quad t > 0,$$

with C_α independent of ε, and the proof is finished. ☐

Remark 4.6. Fix $\varepsilon \geq 0$. Let $\pi_{\alpha,\varepsilon}$ be as in (4.8) with $T_{\alpha,\varepsilon}(t)$ in place of $T_\alpha(t)$ (so $\pi_{\alpha,0} := \pi_\alpha$). The correspondence $\varphi \longmapsto \pi_\alpha(\varphi e_\varepsilon)$ from $\mathcal{T}^{(\alpha)}(t^\alpha)$ into $\mathcal{B}(X)$ is an algebra homomorphism since we have that $(\varphi * g)e_\varepsilon = (\varphi e_\varepsilon) * (g e_\varepsilon)$ for every $\varphi, g \in \mathcal{T}^{(\alpha)}(t^\alpha)$. Moreover, $\pi_{\alpha,\varepsilon}(\varphi) = \pi_\alpha(\varphi e_\varepsilon)$, for all $\varphi \in \mathcal{T}^{(\alpha)}(t^\alpha)$: Take $\varphi \in C_c^\infty[0,\infty)$. By [15, Lemma 5.3],

$$\varphi e_\varepsilon(s) = \sum_{k=0}^\infty \binom{\alpha}{k} \varepsilon^k e^{-\varepsilon s} W^{\alpha-k}\varphi(s), \quad s > 0.$$

Hence, for every $f \in X$, we have

$$\begin{aligned}\pi_{\alpha,\varepsilon}(\varphi)f &= \int_0^\infty W^\alpha \varphi(t) T_{\alpha,\varepsilon}(t) f \, dt \\ &= \int_0^\infty e^{-\varepsilon t} W^\alpha \varphi(t) T_\alpha(t) f \, dt\end{aligned}$$

$$+\sum_{k=1}^{\infty}\binom{\alpha}{k}\int_0^\infty \varepsilon^k e^{-\varepsilon s}W^{\alpha-k}\varphi(s)T_\alpha(s)f\,ds$$
$$=\int_0^\infty\left(\sum_{k=0}^{\infty}\binom{\alpha}{k}\varepsilon^k e^{-\varepsilon s}W^{\alpha-k}\varphi(s)\right)T_\alpha(s)f\,ds=\pi_\alpha(\varphi e_\varepsilon)f.$$

By the density of $C_c^\infty[0,\infty)$ in $\mathcal{T}^{(\alpha)}(t^\alpha)$ we conclude the result.

5. Proof of Theorem 1.1

Proof of Theorem 1.1. First we show that a solution to (1.5) is given by formula (1.6), that is to say, by $u(z)=\pi_\alpha(b^{\sigma,z})f$, for $f\in X$ and $z\in S_{\pi/4}$.

In fact, by Proposition 3.2 (ii),

$$\frac{\partial^2}{\partial z^2}\,b^{\sigma,z}+\frac{1-2\sigma}{z}\,\frac{\partial}{\partial z}\,b^{\sigma,z}=\frac{\partial}{\partial t}\,b^{\sigma,z}$$

in $\mathcal{T}^{(\alpha)}(t^\alpha)$, whence we immediately obtain

$$\frac{\partial^2}{\partial z^2}\,\pi_\alpha(b^{\sigma,z})f+\frac{1-2\sigma}{z}\,\frac{\partial}{\partial z}\,\pi_\alpha(b^{\sigma,z})f=\pi_\alpha\left(\frac{\partial}{\partial t}\,b^{\sigma,z}\right)f,$$

for every $z\in S_{\pi/4}$ and $f\in X$. On the other hand, one has

$$\pi_\alpha\left(\frac{\partial}{\partial t}\,b^{\sigma,z}\right)f=-A\pi_\alpha(b^{\sigma,z})f$$

by (4.9), so $u(z)=\pi_\alpha(b^{\sigma,z})f$ satisfies the differential equation of (1.5). Next, we show that $\pi_\alpha(b^{\sigma,z})f$ satisfies the boundary condition in (1.5).

By using the identity $\dfrac{1}{\Gamma(\alpha)}\displaystyle\int_0^t (t-s)^{\alpha-1}\,ds=\dfrac{t^\alpha}{\Gamma(\alpha+1)}$ and Fubini's theorem one gets

$$\int_0^\infty W^\alpha b^{\sigma,z}(t)\,\frac{t^\alpha}{\Gamma(\alpha+1)}\;dt=\int_0^\infty b^{\sigma,z}(t)\;dt=1,\quad z\in S_{\pi/4}.$$

Take for a moment $f\in\mathcal{D}(A)$. By applying the identity above and (4.3) with $n\ge\alpha$ one obtains

$$\begin{aligned}u(z)-f&=\int_0^\infty W^n(b^{\sigma,z})(t)\left(T_n(t)-\frac{t^n}{\Gamma(n+1)}\right)f\,dt\\&=\int_0^\infty(-1)^n(b^{\sigma,z})^{(n)}(t)T_{n+1}(t)Af\,dt,\end{aligned}$$

where $T_{n+1}(t)=\displaystyle\int_0^\infty (t-s)^{n-\alpha-1}T_{\alpha+1}(s)Af\,\frac{ds}{\Gamma(n-\alpha)}$.

Put $\delta=\Re(z^2)$. By Proposition 3.2(i) and Lemma 3.1(i), one gets

$$\begin{aligned}\|u(z)-f\|&\le\int_0^{\sqrt{\delta}}|(b^{\sigma,z})^{(n)}(t)|\,\|T_{n+1}(t)Af\|\,dt\\&\quad+\int_{\sqrt{\delta}}^\infty|(b^{\sigma,z})^{(n)}(t)|\,\|T_n(t)-\frac{t^n}{n!}\|\,\|f\|\,dt\\&\le C\|b^{\sigma,z}\|_{(n)}\|Af\|\,\sqrt{\delta}\\&\quad+(C+(1/n!))\|f\|\sum_{j=0}^n\frac{|d^\sigma_{j,n}|}{4^{j+\sigma}\Gamma(\sigma)}\,|z|^{2(j+\sigma)}\int_{\sqrt{\delta}}^\infty\frac{e^{-\Re z^2/4t}}{t^{1+\sigma+j}}\,dt\\&\le C_{\sigma,f}\left(\frac{|z^2|}{\Re(z^2)}\right)^{n+\sigma}\sqrt{\delta}\\&\quad+\sum_{j=0}^n C_{j,n}\left(\frac{|z^2|}{\Re(z^2)}\right)^{\sigma+j}\int_{1/\sqrt{\delta}}^\infty\frac{e^{-1/4s}}{s^{1+\sigma+j}}\,ds.\end{aligned}$$

Whence, for f in $\mathcal{D}(A)$, $\lim_{z\to 0} u(z) = f$ uniformly in $S_{\pi/4-\eta}$ for each fixed η such that $0 \le \eta < \pi/4$. Since $\|u(z)\| = \|\pi_\alpha(b^{\sigma,z})f\| \le C_{n,\sigma}\left(\frac{|z^2|}{\Re(z^2)}\right)^{n+\sigma}\|f\|$ for all $f \in X$ and $\mathcal{D}(A)$ is dense in X one obtains that $\lim_{z\to 0} u(z) = f$ uniformly in $S_{\pi/4-\eta}$, for every $f \in X$. This proves the first part of the theorem.

We now prove the second equality of (1.7). Take $n = [\alpha]+1$. Since $T_\alpha(t)$ is a globally tempered α-times integrated semigroup the family $T_n(t)$ given by

$$T_n(t)f := \Gamma(n-\alpha)^{-1}\int_0^t (s-t)^{n-\alpha-1}T_\alpha(s)f\,ds, \quad f \in X,$$

is a globally tempered (of growth n) n-times integrated semigroup as well. Hence, for $f \in X$ and $z \in S_{\pi/4}$,

$$u(z) = (-1)^n\frac{z^{2\sigma}}{4^\sigma\Gamma(\sigma)}\int_0^\infty \frac{d^n}{dt^n}\left(\frac{e^{-z^2/(4t)}}{t^{1+\sigma}}\right)T_n(t)f\,dt,$$

and therefore

$$u'(z) = \frac{(-1)^n z^{2\sigma-1}}{4^\sigma\Gamma(\sigma)}\int_0^\infty \frac{d^n}{dt^n}\left(\left(2\sigma - \frac{z^2}{2t}\right)\frac{e^{-z^2/(4t)}}{t^{1+\sigma}}\right)T_n(t)f\,dt. \tag{5.1}$$

On the other hand,

$$\int_0^\infty \frac{d^n}{dt^n}\left(\left(2\sigma - \frac{z^2}{2t}\right)\frac{e^{-z^2/(4t)}}{t^{1+\sigma}}\right)\frac{t^n}{n!}\,dt = \int_0^\infty\left(\left(2\sigma - \frac{z^2}{2t}\right)\frac{e^{-z^2/(4t)}}{t^{1+\sigma}}\right)dt = 0$$

That the second integral vanishes can be seen by making the change of variable $r = z^2/4t$ for $z > 0$; then the integral is null for every $z \in S_{\pi/4}$ by the analytic continuation principle.

According to the two preceding remarks we have, for every $f \in X$,

$$z^{1-2\sigma}u'(z) = \frac{(-1)^n}{4^\sigma\Gamma(\sigma)}\int_0^\infty \frac{d^n}{dt^n}\left(\left(2\sigma - \frac{z^2}{2t}\right)\frac{e^{-z^2/(4t)}}{t^{1+\sigma}}\right)\left(T_n(t)f - \frac{t^n}{n!}f\right)dt$$

which, once having used the product derivation rule and put $a = -z^2/4$, gives us

$$\begin{aligned} z^{1-2\sigma}u'(z) &= \sum_{k=0}^n\sum_{m=0}^k D^\sigma_{m,k,n}\,a^{m+1}\int_0^\infty \frac{e^{a/t}}{t^{\sigma+m+n+2}}\left(T_n(t)f - \frac{t^n}{n!}f\right)dt \\ &\quad + \sum_{m=0}^n C^\sigma_{m,n,n}\,a^m\int_0^\infty \frac{e^{a/t}}{t^{\sigma+m+n+1}}\left(T_n(t)f - \frac{t^n}{n!}f\right)dt, \end{aligned}$$

with

$$C^\sigma_{0,n,n} = \frac{2\sigma(-1)^n}{4^\sigma\Gamma(\sigma)}\,d^\sigma_{0,n} = \frac{2\sigma}{4^\sigma\Gamma(\sigma)}\,c_{0,n,\sigma} = \frac{2\sigma}{4^\sigma\Gamma(\sigma)}\,\frac{\Gamma(\sigma+n+1)}{\Gamma(\sigma+1)},$$

where $d^\sigma_{0,n}$ and $c_{0,n,\sigma}$ are given by (3.4) and (3.3) respectively.

Take now $f \in \mathcal{D}(A)$. We are going to see that the additive terms in the expression above of $z^{1-2\sigma}u'(z)$ tend to 0, as $a \to 0$, with the only exception of the term with constant $C^\sigma_{0,n,n}$. This last term will allow us to obtain the fractional power $(-A)^\sigma$.

Actually, for $1 \le j \le n+1$,

$$\begin{aligned} &|a|^j\int_0^\infty \frac{e^{(\Re a)/t}}{t^{\sigma+j+n+1}}\left\|T_n(t)f - \frac{t^n}{n!}f\right\|dt \\ &\qquad\qquad \le |a|^j\int_0^\infty \frac{e^{(\Re a)/t}}{t^{\sigma+j+n+1}}\,\|T_{n+1}(t)\|\|Af\|\,dt \\ &\qquad\qquad \le C|a|^j\int_0^\infty \frac{e^{(\Re a)/t}}{t^{\sigma+j}}\,dt \end{aligned}$$

$$= C\left(\frac{|a|}{\Re a}\right)^j \left(\int_0^\infty \frac{e^{1/s}}{s^{\sigma+j}}\,ds\right)(\Re a)^{1-\sigma},$$

where we have applied (4.3) in the first inequality. Moreover,

$$\lim_{a\to 0}\int_0^\infty \frac{e^{a/t}}{t^{\sigma+n+1}}\Big(T_n(t)f-\frac{t^n}{n!}f\Big)\,dt = \int_0^\infty \frac{(T_n(t)f-(t^n/n!)f)}{t^{\sigma+n+1}}\,dt$$

by the Dominated Convergence Theorem (use again (4.3) near the origin).

Then, making z converging to 0 through subsectors of $S_{\pi/4}$, and applying Theorem 4.1, we obtain

$$\begin{aligned}\lim_{z\to 0} z^{1-2\sigma}u'(z) &= \frac{2\sigma}{4^\sigma\Gamma(\sigma)}\frac{\Gamma(\sigma+n+1)}{\Gamma(\sigma+1)}\frac{\Gamma(-\sigma)\Gamma(\sigma+1)}{\Gamma(\sigma+n+1)}(-A)^\sigma f\\ &= \frac{2\sigma\Gamma(-\sigma)}{4^\sigma\Gamma(\sigma)}(-A)^\sigma f,\end{aligned}$$

as we wanted to prove.

Next, we are going to see that the solution $u(z)$ above satisfies formula (1.8). By Lemma 3.5, $B^{\sigma,z}=b^{\sigma,z}*h^\sigma$, $z\in S_{\pi/4}$. Thus we have $B^{\sigma,z}e_\varepsilon=(b^{\sigma,z}e_\varepsilon)*(h^\sigma e_\varepsilon)$ in $\mathcal{T}^{(\alpha)}(t^\alpha)$ for every $\varepsilon>0$. Moreover,

$$(\varepsilon-A)^{-\sigma}=\pi_\alpha(e_\varepsilon h^\sigma),\quad \varepsilon>0; \tag{5.2}$$

see [26, Remark 4.10]. Then, by Proposition 3.2(iii), (4.6), (5.2) and Lemma 3.5, we get for $f\in\mathcal{D}(A)$ that

$$\begin{aligned}u(z)=\pi_\alpha(b^{\sigma,z})f &= \lim_{\varepsilon\to 0^+}\pi_\alpha(b^{\sigma,z}e_\varepsilon)(\varepsilon-A)^{-\sigma}(-A)^\sigma f\\ &= \lim_{\varepsilon\to 0^+}\pi_\alpha(b^{\sigma,z}e_\varepsilon)\pi_\alpha(h^\sigma e_\varepsilon)(-A)^\sigma f=\lim_{\varepsilon\to 0^+}\pi_\alpha(B^{\sigma,z}e_\varepsilon)(-A)^\sigma f,\end{aligned}$$

uniformly in z running over compact subsets of $S_{\pi/4}$. Thus (1.8) is proved.

Formula (1.9) is rightly obtained from the above. In fact, that formula can be written as

$$u(z)-f=\pi_\alpha(B^{\sigma,z}-h^\sigma)(-A)^\sigma f. \tag{5.3}$$

To see that (5.3) holds one only needs to apply the limit above and Proposition 3.6 (ii):

$$\begin{aligned}u(z)-f &= \lim_{\varepsilon\to 0^+}\big(\pi_\alpha(B^{\sigma,z}e_\varepsilon)(-A)^\sigma f-\pi_\alpha(h^\sigma e_\varepsilon)(-A)^\sigma f\big)\\ &= \lim_{\varepsilon\to 0^+}\pi_\alpha\left((B^{\sigma,z}-h^\sigma)e_\varepsilon\right)(-A)^\sigma f\\ &= \pi_\alpha(B^{\sigma,z}-h^\sigma)(-A)^\sigma f.\end{aligned}$$

It remains to prove the characterization of the fractional power $(-A)^\sigma$ which is indicated in the first equality of formula (1.7). Notice that

$$\begin{aligned}\frac{z^{-2\sigma}}{\Gamma(\sigma)}\int_0^\infty \frac{d^n}{dt^n}\left(\frac{e^{-z^2/(4t)}-1}{t^{1-\sigma}}\right)\frac{t^n}{n!}\,dt &= \frac{z^{-2\sigma}}{\Gamma(\sigma)}\int_0^\infty \frac{e^{-z^2/(4t)}-1}{t^{1-\sigma}}\,dt\\ &= \frac{4^{-\sigma}}{\Gamma(\sigma)}\int_0^\infty r^{\sigma-1}(e^{-1/r}-1)\,dr\\ &= -\frac{4^{-\sigma}}{\Gamma(\sigma)}\int_0^\infty\int_0^s e^{-u}\,du\,s^{-(1+\sigma)}\,ds\\ &= -\frac{4^{-\sigma}}{\sigma\Gamma(\sigma)}\int_0^\infty u^{-\sigma}e^{-u}\,du=\frac{\Gamma(-\sigma)}{4^\sigma\Gamma(\sigma)}=c_\sigma.\end{aligned}$$

Hence,

$$\begin{aligned}&z^{-2\sigma}(u(z)-f)-c_\sigma(-A)^\sigma f\\ &\quad=\frac{z^{-2\sigma}}{\Gamma(\sigma)}\int_0^\infty \frac{d^n}{dt^n}\left(\frac{e^{-z^2/(4t)}-1}{t^{1-\sigma}}\right)\left(T_n(t)-\frac{t^n}{n!}\right)(-A)^\sigma f\,dt\end{aligned}$$

$$
\begin{aligned}
&= \frac{z^{-2\sigma}}{\Gamma(\sigma)} \int_0^{\sqrt{\Re(z^2)}} \frac{d^n}{dt^n}\left(\frac{e^{-z^2/(4t)}-1}{t^{1-\sigma}}\right) T_{n+1}(t)A(-A)^\sigma f\,dt\\
&+ \frac{z^{-2\sigma}}{\Gamma(\sigma)} \int_{\sqrt{\Re(z^2)}}^{\infty} \frac{d^n}{dt^n}\left(\frac{e^{-z^2/(4t)}-1}{t^{1-\sigma}}\right) \left(T_n(t)-\frac{t^n}{n!}\right)(-A)^\sigma f\,dt.
\end{aligned}
$$

Thus

$$
\begin{aligned}
&\|z^{-2\sigma}(u(z)-f)-c_\sigma(-A)^\sigma f\|\\
&\le C\frac{|z|^{-2\sigma}}{\Gamma(\sigma)}\sqrt{\Re(z^2)}\,\|A(-A)^\sigma f\| \int_0^{\sqrt{\Re(z^2)}} \left|\frac{d^n}{dt^n}\left(\frac{e^{-z^2/(4t)}-1}{t^{1-\sigma}}\right)\right| t^n dt\\
&+ C\frac{|z|^{-2\sigma}}{\Gamma(\sigma)} \int_{\sqrt{\Re(z^2)}}^{\infty} \left|\frac{d^n}{dt^n}\left(\frac{e^{-z^2/(4t)}-1}{t^{1-\sigma}}\right)\right| t^n\,dt.
\end{aligned}
$$

Then the first member of the right hand of the inequality is bounded by

$$
\begin{aligned}
C|z|^{-2\sigma}\sqrt{\Re(z^2)}\,\|B^{\sigma,z}-h^\sigma\|_{(n)} &\le C_{n,\sigma}\left(|z^2|^{-\sigma}\sqrt{\Re(z^2)}\right)\frac{|z|^{2n}}{(\Re(z^2))^{n-\sigma}}\\
&\le C_{n,\sigma}\left(\frac{|z^2|}{\Re(z^2)}\right)^n \sqrt{\Re(z^2)},
\end{aligned}
$$

by (3.6), whereas the second member is bounded by

$$
\begin{aligned}
&C|z|^{-2\sigma}\int_{\sqrt{\Re(z^2)}}^{\infty}\left(\frac{1-e^{-\Re(z^2)/(4t)}}{t^{1-\sigma}}\right)dt\\
&\qquad+\sum_{k=1}^{n}\sum_{m=1}^{k} C_{m,k}|z|^{-2\sigma}\left|-\frac{z^2}{4}\right|^m \int_{\sqrt{\Re(z^2)}}^{\infty}\frac{e^{-\Re(z^2)/(4t)}}{t^{m+1-\sigma}}\,dt\\
&= C\left(\frac{\Re(z^2)}{|z^2|}\right)^{\sigma}\int_{1/\sqrt{\Re(z^2)}}^{\infty}\left(\frac{1-e^{-1/(4s)}}{s^{1-\sigma}}\right)ds\\
&\qquad+\sum_{k=1}^{n}\sum_{m=1}^{k} C_{m,k}\left(\frac{|z^2|}{\Re(z^2)}\right)^{m-\sigma}\int_{1/\sqrt{\Re(z^2)}}^{\infty}\frac{e^{-1/(4s)}}{s^{m+1-\sigma}}\,ds.
\end{aligned}
$$

Putting all the preceding estimates together one obtains that, as it was claimed, $\lim_{z\to 0} z^{-2\sigma}(u(z)-f)=c_\sigma(-A)^\sigma f$, $f\in\mathcal{D}(A)$, through subsectors of $S_{\pi/4}$. We have proved all the formulas of the theorem.

Finally, the boundedness of the solution u on subsectors of $S_{\pi/4}$ is easy to show. Take $n>\alpha$. Since A is also the generator of the integrated semigroup $T_n(t)$ defined in (4.2), it follows by (1.6) applied to $T_n(t)$ and Proposition 3.2 (i) that

$$
\|u(z)\| \le C_n\frac{|z^{2\sigma}|}{4^\sigma\Gamma(\sigma)}\|f\|\int_0^\infty \frac{d^n}{dt^n}\left(\frac{e^{-z^2/(4t)}}{t^{1+\sigma}}\right)t^n\,dt \le C_{n,\sigma}\left(\frac{|z^2|}{\Re(z^2)}\right)^{n+\sigma}\|f\|,\quad z\in S_{\pi/4},
$$

as we wanted to show. □

Acknowledgments. The third author wishes to thank the Departamento de Matemáticas at Universidad de Zaragoza for its kind hospitality during several visits.

References

[1] W. Arendt, C. J. K. Batty, M. Hieber and F. Neubrander, *Vector-valued Laplace Transforms and Cauchy Problems*, Monographs in Math. **96**, Birkhäuser, Basel, 2001.

[2] W. Arendt and H. Kellerman, Integrated solutions of Volterra integro-differential equations and applications, in: *Integrodiff. Eq., Proc. Conf. Trento 1987*, (G. Da Prato and M. Iannelli, eds.), Pitman Res. Notes Math. **190**, Longman, Harlow, 1987, pp. 21-51.

[3] L. Caffarelli, S. Salsa and L. Silvestre, Regularity estimates for the solution and the free boundary of the obstacle problem for the fractional Laplacian. *Invent. Math.* **171** (2008), 425–461.
[4] L. Caffarelli and L. Silvestre, An extension problem related to the fractional Laplacian, *Comm. Partial Differential Equations* **32** (2007), 1245–1260.
[5] L. Caffarelli and A. Vasseur, Drift diffusion equations with fractional diffusion and the quasi-geostrophic equation, *Ann. of Math. (2)* **171** (2010), 1903–1930.
[6] G. Carron, T. Coulhon and E.-M. Ouhabaz, Gaussian estimates and L^p-boundedness of Riesz means, *J. Evol. Equ.* **2** (2002), 299–317.
[7] S-Y. A. Chang, M. M. González, Fractional Laplacian in conformal geometry, *Adv. Math.* **226** (2011), 1410–1432.
[8] E. B. Davies, L^p spectral theory and L^1 analyticity *J. London Math.Soc.* **52** (1995), 177–184.
[9] E. B. Davies, Uniformly elliptic operators with measurable coefficients *J. Funct. Anal.* **132** (1995), 141–169.
[10] N. Dunford and J. T. Schwarz, *Linear Operators, Part III: Spectral Operators*, Pure and Applied Math. **VII**, Wiley-Interscience, New York, 1971.
[11] O. El-Mennaoui, *Traces de semi-groupes holomorphes singuliers A l'origine et comportement asymptotique*, Thése, Besancon 1992.
[12] O. El-Mennaoui and V.Keyantuo, Trace theorems for holomorphic semigroups and the second order Cauchy problem, *Proc. Amer. Math. Soc.* **124** (1996), 1445–1458.
[13] J. E. Galé and P. J. Miana, One parameter groups of regular quasimultipliers, *J. Funct. Anal.* **237** (2006), 1–53.
[14] J. E. Galé, P. J. Miana and J. J. Royo, Estimates of the Laplace transform on convolution Sobolev algebras, *J. Approx. Theory* **164** (2012), 162–178.
[15] J. E. Galé and T. Pytlik, Functional calculus for infinitesimal generators of holomorphic semigroups, *J. Funct. Anal.* **150** (1997), 307–355.
[16] J. A. Goldstein, *Semigroups of Linear Operators and Applications*, Oxford University Press, New York, 1985.
[17] M. Hieber, Integrated semigroups and differential operators on L^p spaces, *Math. Ann.* **291** (1991), 1–16.
[18] M. Hieber, Integrated semigroups and the Cauchy problem for systems in L^p spaces, *J. Math. Anal. Appl.* **162** (1991), 300–308.
[19] M. Hieber, Laplace transforms and α-times integrated semigroups, *Forum Math.* **3** (1991), 595–612.
[20] M. Hieber, L^p spectra of pseudodifferential operators generating integrated semigroups *Trans. Amer. Math. Soc.* **347** (1995), 4023–4035.
[21] M. Hieber, Spectral theory and Cauchy problems on L^p spaces, *Math. Z.* **216** (1993), 613–628.
[22] Y. Hu and G. Kallianpur, Schrödinger equations with fractional Laplacians, *Appl. Math. Optim.* **42** (2000), 281–290.
[23] V. Keyantuo, Integrated semigroups and related partial differential equations, *J. Math. Anal. Appl.* **212** (1997), 135–153.
[24] N. N. Lebedev, *Special Functions and Their Applications*, Revised English edition. Translated and edited by Richard A. Silverman, Prentice-Hall, Inc., Englewood Cliffs, NJ, 1965.
[25] C. Martínez, M. Sanz and L. Marco, Fractional powers of operators, *J. Math. Soc. Japan*, **40** (1988), 331–347.
[26] P. J. Miana, α-times integrated semigroups and fractional derivation, *Forum Math.* **14** (2002), 23–46.
[27] D. Müller, Functional calculus for Lie groups and wave propagation, Doc. Math., International Congress Math., Berlin, 1998.
[28] D. Müller and E. M. Stein, L^p estimates for the wave equation on $\mathbb{H}^n$, *Rev. Matem. Iberoamericana* **15** (1999), 293–334.
[29] J. Randall, The heat kernel for generalized Heisenberg groups, *J. Geom. Anal.* **6** (1996), 287–316.
[30] L. Roncal and P. R. Stinga, Fractional Laplacian on the torus, preprint 2012.
[31] P. R. Stinga and J. L. Torrea, Extension problem and Harnack's inequality for some fractional operators, *Comm. Partial Differential Equations* **35** (2010), 2092–2122.
[32] P. R. Stinga and C. Zhang, Harnack's inequalities for fractional nonlocal equations, *Discrete Contin. Dyn. Syst.* (2012), to appear; arXiv: 1203 . 1518v2.
[33] E. Taflin, Analytic linearization of the Korteweg-De Vries equation, *Pacific J. Math.* **108** (1983), 203–220.
[34] O. Vallée and M. Soares, *Airy Functions and Applications to Physics*, World Scientific, Nerw Jersey, 2004.
[35] J. M. A. M. van Neerven and B. Straub, On the existence and growth of mild solutions of the abstract Cauchy problem for operators with polynomially bounded resolvent, *Houston J. Math.* **24** (1998), 137–171.
[36] K. Yosida, *Functional Analysis*, Springer-Verlag, New York, 1978.
[37] G. M. Zaslavsky, Chaos, fractional kinetics, and anomalous transport, *Phys. Rep.* **371** (2002), 461–580.

Departamento de Matemáticas e I. U. M. A. Universidad de Zaragoza, 50009 Zaragoza, Spain
E-mail address: `gale@unizar.es, pjmiana@unizar.es`

Departamento de Matemáticas y Computación, Universidad de La Rioja, 26004 Logroño, Spain
E-mail address: `pablo-raul.stinga@unirioja.es`